

*eyes* moderate, oval, somewhat convex. *Thorax* narrowed to base; *pronotum* convex, rounded in front and at sides, *pro-mesonotal* suture distinct; *meso-epinotal* suture wanting; *epinotum* unarmed, gradually sloping to base, no marked angle between dorsal surface and declivity. Node of *pedicel* simple, unarmed, convex, rounded anteriorly and at sides; *gaster* narrowed to base, the constriction between the first and second segments very marked, the puncturation of the last segments not nearly so coarse as that of the rest of the body.

*Long.* 11 mm.

This distinct species comes nearest to *tesserinoda*, but besides being considerably larger the sculpture is entirely different.

Described from two workers. Dohnavur, Tinnevely District, S. India, 30. ix. 38.

Type in B.M. (N.H.).

I have named this species in honour of Mr. G. M. Henry, who captured the ants of this expedition.

4. *Euponera* (*Mesoponera*) *melanaria* Emery, Ann. Soc. Ent. France, lxii. 260, note (1893).

♂ (undescribed). Brownish yellow, eyes black, antennæ except scape and first joint of funiculus dark brown, covered with long outstanding and short decumbent yellow hairs, shining, very finely punctured.

*Head* about as long as broad, rounded behind eyes, posterior border slightly emarginate, with a narrow raised margin; *mandibles* short, reduced, bluntly pointed; *clypeus* convex, rounded anteriorly; *clypeal foveæ* round, deep; *suture* between clypeus and frontal area distinct; *frontal area* large, flat, triangular, finely but distinctly defined; *frontal carinæ* very short, round; *antennæ* long, 13-jointed; *scape* short, slightly longer than broad; *funiculus* with 1st joint very short, transverse, considerably shorter and somewhat narrower than scape; the rest of the joints longer than broad, of about equal breadth, last joint slightly longer than the preceding; *eyes* large, kidney-shaped, situated on each side of the head; *ocelli* moderate, oval, placed on vertex of head. *Thorax* convex, narrowed in front and behind, broadest just before insertion of the wings; *pronotum* transverse, furnished with a short, finely margined neck, dorsal surface convex, sides encircling the mesonotum; *meso-*