notum very convex, broad oval, sides and front rounded. a short pointed carina extends from centre of anterior border ending in a rather broad short furrow; mayrian furrows wanting; parapsidal furrows narrow but distinct; the mesonotum and scutellum are divided by a short transverse furrow; scutellum convex and prominent; metanotum narrow, transverse, somewhat raised; epinotum long, sloping, no angle between dorsal surface and declivity. Petiole furnished with a rather high node, straight posteriorly, sloping anteriorly, armed beneath with a projection toothed at apex and base; gaster short, oval, constriction between post-petiole and second segment well marked; pygidium armed with a long, sharp, curved cerci well developed; genitalia prominent; stipites large, curved, bluntly pointed; sagitæ sharply Wings iridescent, veins and pterostigma light brown; two discoidal, two cubital cells present; radial cell long, closed, slightly appendiculate.

Long. 6.3 mm.

Described from a single specimen, Nadungayam, 200 ft., Malabar, S. India, 16-22 ix. 38. Male type in B.M. N.H.).

5. Cryptopone testacea Mots., Bull. Soc. Nat. Mosco, xxxvi. p. 15 (1863), Q; Emery, Ann. Soc. Ent. France, 1893, p. 240, \mathsepsilon .

Seven winged QQ. Nadungayam, 200 ft., Malabar, S. India, 16-22. ix. 38.

Distribution.—Ceylon.

 \mathcal{F} (undescribed). Yellow, eyes black, pubescence, sculpture and colour much as in \mathcal{P} .

Head across eyes about as broad as long, slightly narrowed posteriorly, posterior angles rounded; mandibles small, much reduced, pointed at apex; clypeus convex, anterior border rounded; clypeal foveæ very small, round; cheeks very short; frontal carinæ very short, round, with a small shallow hollow between; frontal furrow short; antennæ missing; eyes large, somewhat kidney-shaped, situated at sides of head, anterior border very close to posterior border of clypeus; ocelli round, prominent, not close together. Thorax not broader than head together with eyes, narrowed posteriorly; pronotum transverse, margined and furnished with a neck anteriorly, sinuate on each side; mesonotum convex, oval, slightly broader than long,