

Note.—In Bingham's table (*l. c.*) he gives for *A. yerburyi* Forel: "Pro-, meso- and metanotum punctate, not striate," and in the description, "Head . . . faintly longitudinally striate. . . . The meso- and mesonotum transversely striate"!

Subfamily **Pseudomyrininæ.**

8. *Sima rufonigra* Jerd., Madras Journ. Lit. Sci. xvii.* p. 111 (1851).

One del. ♀, 2 ♀♀. Trenmalat, 500–800 ft., Travancore, S. India (♀); Dohnavur, 350 ft., Tinnevelly District, S. India, i. x. 38 (♀♀).

Distribution.—Continental India, Ceylon, Sumatra.

9. *Tetraponera nigra* Jerd., Madras Journ. Lit. Sci. xvii. p. 112 (1851).

One del. ♀. Dohnavur, 350 ft., Tinnevelly District, S. India, i. x. 38.

Distribution.—Continental India, Ceylon.

Subfamily **Myrmicinæ.**

10. *Pheidole* (*Pheidole*), sp. ?

Ten ♀♀. Dohnavur, 350 ft., Tinnevelly District, S. India, 30. ix. 38.

It is quite impossible to name the workers of *Pheidole* species unless a soldier be present also.

11. *Myrmicaria brunnea* Saunders, Trans. Ent. Soc. Lond. iii. p. 57 (1841).

Eight ♀♀. Nadungayam, 200 ft., Malabar, S. India, 16–22. ix. 38, and Dohnavur, 350 ft., Tinnevelly District, S. India, 30. ix. 38.

Distribution.—India, Ceylon, Sumatra.

12. *Crematogaster* (*Acrocælia*) *brunnea* Smith, subsp. *subnuda* Mayr, Verh. Zool. Bot. Ges. Wien, xxviii. p. 682 (1878).

Three ♀♀. Dohnavur, 350 ft., Tinnevelly District, S. India, 30. ix. 38.

Distribution.—India, Assam, Burma.

* In the 'Genera Insectorum' the volume is incorrectly given as "15."