Crematogaster (Acrocælia) rothneyi Mayr, Verh. Zool. Bot. Ges. Wien, xxviii. p. 685 (1878).

One  $\mbox{$ \xi$}$ . Dohnavur, 350 ft., Tinnevelly District, S. India, 1. x. 38.

Distribution.—India.

14. Monomorium (Holcomyrmex) glabrum André, Spec. Hym. Europe, ii. p. 345 (1883).

Four \( \beta \beta \). Dohnavur, 350 ft., Tinnevelly District, S. India, 1. x. 38.

Distribution.—The whole of India, Ceylon and Burma.

15. Monomorium (Holcomyrmex) scabriceps Mayr, Verh. Zool. Bot. Ges. Wien, xxviii. p. 672 (1878).

One  $\normalfont{Q}$ . Dohnavur, Tinnevelly District, 350 ft., S. India, 8. x. 38.

Distribution.—India.

16. Meranoplus levis, sp. n.

\$\times\$. Head, thorax, antennæ, legs, first joint of pedicel, and anterior margin of gaster very narrowly, yellowish red, eyes and teeth of mandibles black, club of antennæ brown, post-petiole reddish brown, bordered with black, gaster very dark brown, almost black. Head, thorax and petiole mostly smooth, shining, clothed with longer and shorter outstanding yellow hairs, thicker on the gaster.

Head slightly longer than broad, narrowed in front, sides rounded, sinuate before posterior angles, posterior angles pointed, posterior border truncate, vertex, with a few faint longitudinal striæ; mandibles triangular, closely longitudinally striate, armed with four sharp teeth; clypeus flat, with a few longitudinal striæ, a narrow carina on each side ends in a small sharp tooth at corners of anterior border, which is margined and truncate; frontal area indistinct; frontal carinæ pointed, widely separated, continued to form a margin above the large wide, shallow, smooth antennal grooves; antennæ fairly long, 9-jointed, with a 3-jointed club; cheeks with several raised longitudinal ridges; eyes moderate, round-oval, prominent.

Thorax transverse; pro-mesonotal shield transverse, convex, covered with a number of faint oblique longitudinal striæ, narrowly margined anteriorly, sinuate before the short, pointed, anterior angles, posterior angles armed with a rather long pointed spine, posterior border margined; epinotum concave, margined at sides, armed with two long