thin sharp spines; petiole flat, transverse, margined; post-petiole globose, margined, slightly narrowed posteriorly, covered with a network of raised ridges; gaster heart-shaped, anterior border excised and narrowly margined, sculpture of anterior portion consisting of finely raised longitudinal ridges and very fine punctures.

Long. 4.3 mm.

Described from two workers, Dohnavur, 350 ft., Tinnevelly District, S. India, 1. x. 38.

Type in B.M. (N.H.).

This very distinct species is chiefly distinguished by its mostly smooth surface.

There are four specimens of this species in the B.M. from Ceylon: "S. P. Hambantota, T. B. F., 6 & 7 Feb. '09. Ceylon, T. B. Fletcher, 1909—137."

17. Meranoplus carinatus, sp. n.

Q. Head, thorax and legs black, cheeks yellow above, petiole and post-petiole yellowish brown, gaster brownish yellow with dark markings, clothed with longer and shorter outstanding curved yellow hairs, thicker on gaster. Sculpture of dorsal surface of head and thorax consisting of longitudinal raised ridges, or carinæ, which form a network and enclose larger and smaller, round, shallow punctures.

Head triangular, narrowed in front, broadest behind, posterior angles round, prominent, posterior border slightly but widely excised; mandibles massive, triangular, covered with numerous fine raised ridges, armed with four strong teeth; clypeus concave before antérior border, which is truncate, smooth and shining, with scattered punctures, thicker at the sides; frontal area transverse, triangular, smooth and shining; cheeks with strong longitudinal raised ridges; antennal grooves wide, deep, smooth and shining anteriorly, striate posteriorly; antennæ missing; eyes large, convex, round-oval; ocelli moderate. Thorax triangular, broadest across anterior angles, dorsal surface flat, anterior angles pointed; suture between mesonotum and scutellum narrow but deep; scutellum transverse, triangular; epinotum armed with two short, broad, but pointed teeth, dorsal surface shorter than declivity, longitudinally striate, declivity somewhat concave, very smooth and shining. Petiole with a rather high node, bidentate at apex, anterior surface slanting.