

22. *Camponotus (Tanæmyrmex) barbatus* Rog., sub. sp. *taylori* Forel, Journ. Bombay Nat. Hist. Soc. vii. p. 241 (1892).

One ♀. Tenmalai, 500–800 ft., Travancore, S. India, 11–17. x. 38.

*Distribution*.—India.

23. *Camponotus (Tanæmyrmex) variegatus* Smith, Cat. Hym. Brit. Mus. vi. p. 19 (1858).

Two large ♂♂. Dohnavur, 350 ft., Tinnevelly District, S. India, 1. x. 38.

*Distribution*.—Ceylon, Burma, Singapore.

24. *Camponotus (Tanæmyrmex) compressus* F., Mant. Ins. i. p. 307 (1787).

Thirty-two (4 ♂ and ♀♀). Shencotta, S. India, 11. x. 38; Dohnavur, 350 ft., Tinnevelly District, S. India, 1. x. 38.

*Distribution*.—India, Ceylon.

25. *Camponotus (Myrmosericus) rufoglaucus* Jerdon, Madras Journ. Lit. Sci. xvii. p. 124 (1851).

Thirty ♀♀. Dohnavur, 350 ft., Tinnevelly District, S. India, 8. x. 38.

*Distribution*.—India, Ceylon, Assam, Burma.

26. *Camponotus (Myrmosericus) rufoglaucus* Jerdon, subsp. *paria* Emery, Ann. Mus. Stor. Nat. Genova, xxvii. p. 513 (1889).

Two ♀♀. Nadungayam, 200 ft., Malabar, S. India, 16–22. ix. 38.

*Distribution*.—India, Ceylon, Assam, Burma, Siam.

27. *Camponotus (Orthonotomyrmex) sericeus* F., Suppl. Ent. Syst. p. 279 (1798).

One del. ♀, 25 ♀♀. Tuticorin, S. India, 18. x. 38. This was probably a young colony, as none of the workers are large enough to be called soldiers. Sixty (4 ♂ and ♀♀). Dohnavur, 350 ft., Tinnevelly District, S. India, 30. ix., 1 & 8. x. 38. Some of these soldiers, and also workers, possess blood-red heads, =var. *sanguiniceps* Donis.

*Distribution*.—Tropical Africa, Egypt, Arabia, India, Indochina, Ceylon.

28. *Camponotus (Orthonotomyrmex ?) puniceps*, sp. n.

♀. Head, thorax, pedicel, coxae and trochanters reddish pink, antennæ and spurs reddish yellow, anterior pair of legs brownish red, posterior pairs of legs and gaster black.