

the ridges and hollows beneath the epinotum are much stronger. Long. 8.2 mm.

CEYLON: S.P. Humbantota, J. B. Fletcher, 6 Oct., 1909, 1 ♀ (type, in coll. British Museum (Nat. Hist.)).

As pointed out in my description of *Camponotus (Orthonotomyrmex?) puniceps*, that species, except for the form of the scale, would fit in the subgenus *Orthonotomyrmex*, and it is not wise to create a new subgenus on this alone. As it comes so close to *C. (O.) fletcheri*, which is undoubtedly an *Orthonotomyrmex*, it would appear best to leave *puniceps* in this subgenus.

REFERENCES.

- ASHMEAD, W. H. 1905. *Can. Ent.*, **37**: 384.
 ———— 1906. *Proc. Ent. Soc., Wash.*, **8**: 31.
 BINGHAM, C. T. 1903. *Faun. Brit. India, Hym.*, **2**: 370, 376.
 DONISTHORPE, H. ST. J. K. 1942. *Ann. Mag. nat. Hist.*, (11) **9**: 458-9.
 EMERY, C. 1895. *Ann. Soc. ent. Fr.*, **63**: 54.
 ———— 1920. *Rev. Zool. Afr.*, **8**: 229-60.
 ———— 1925. *Gen. Ins.*, **183**: 65, 124.
 FOREL, A. 1895. *J. Bombay nat. Hist. Soc.*, **9**: 454.
 ———— 1907. *Ann. Mus. nat. Hung.*, **5**: 31.
 ———— 1912. *Mém. Soc. ent. Belg.*, **20**: 92.
 ———— 1914. *Rev. Suisse Zool.*, **22**: 264, 273.
 SANTSCHI, F. 1921. *Ann. Soc. ent. Belg.*, **61**: 310.

British Museum (Nat. Hist.),
 London, S.W.7.

April 29th, 1942.