

tracted behind pronotum; *pronotum* large, transverse, convex, sides and anterior angles rounded; *prosternum* forming a neck anteriorly; *mesonotum* circular, about as long as broad; *epinotum* longer than broad, narrowed behind, sloping to base, without a distinct angle between dorsal surface and declivity. *Scale of petiole* not very high, longer than broad, narrowed to base, apex bluntly pointed and showing a tendency to being slightly excised; *gaster* short oval, broadest at base, rounded anteriorly, narrowed to apex. *Legs* moderately long. *Long.* 2.5–2.7 mm.

♀. Black, shining, legs, antennae, and mandibles brown, apex of antennae and tarsi pale brownish-yellow, clothed rather thickly with yellowish-white pubescence; whiter and thicker on cheeks, and some short yellow outstanding hairs, especially on mandibles and gaster.

*Head* slightly longer than broad, narrowed anteriorly; sides behind eyes convex and rounded, posterior angles rounded, posterior border distinctly emarginate; *mandibles* massive, triangular, punctured, armed with one rather long curved tooth at apex and a small tooth just before it; the rest of the masticatory border faintly serrate; *clypeus* large, transverse, rather flat, anterior border rounded, *frontal area* not indicated; *eyes* large, round oval, very slightly convex, situated on each side of the head before the centre, touching the lateral borders when seen from above; *ocelli* very small, situated on top of posterior part of head near posterior border; *antennae* long, *scape* extending a little beyond the posterior border of the head, *funiculus* with first two joints distinctly longer than broad, the 1st joint narrower but if anything slightly longer than the second, 3rd joint about as long as broad; 4th–10th slightly longer than broad, last joint very slightly longer than the two preceding taken together. *Thorax* large, massive, somewhat flat above, broadest across anterior angles, narrowed to base; *pronotum*, transverse, narrow, anterior border slightly raised, *prosternum* forming a neck; *mesonotum* large, transverse, flat, rounded anteriorly and posteriorly; *praescutellum* indistinct; *scutellum* large, transverse, flat, narrowed to base; *epinotum* about as long as broad, sides straight, dorsal surface convex, shorter than the declivity, declivity rather abrupt, but angle between it and dorsal surface not marked. *Scale of petiole* hexagonal, not large, anterior border emarginate, sides with shortly pointed angles, space between anterior border and angles excised, sides behind angles straight, narrowed to base. *Gaster* short, round oval, narrowed to apex. *Legs* fairly long. *Wings* iridescent, *pterostigma* and *veins* pale brown, one cubital cell, one discoidal cell, and rather narrow, long, not quite closed radial cell present. *Long.* 3.7–4 mm.

♂. Black, shining, mandibles, legs and antennae pale brown, tarsi and apex of antennae lighter, clothed with fine greyish pubescence and a few short yellow outstanding hairs.

*Head* transverse, narrowed before eyes, sides behind eyes rounded, posterior border truncate; *mandibles* long, narrow, crossed at apex when closed, armed with a sharp narrow tooth at apex, and the indication of a smaller tooth preceding it; *clypeus* transverse, convex, anterior border round and slightly projecting in centre, slightly sinuate on each side; *vertex* of head slightly concave; *eyes* large, oval, prominent, situated on each side of the head in front, close to posterior border of clypeus, but with an evident space between; *ocelli* fairly large, prominent, larger in proportion than those of the female, situated on top of posterior part of head which is slightly raised; *antennae* moderately long, narrow, *scape* extending as far as the median ocellus, all the joints of the *funiculus* longer than broad, first joint shorter than the rest, last joint not quite as long as the two preceding taken together. *Thorax* longer than broad, convex, narrowed in front and behind, broadest across middle of mesonotum; *pronotum* transverse, narrow anteriorly, sides extending round mesonotum to insertion of fore-wing; *prosternum* forming a neck; *mesonotum* transverse, convex, rounded in front and at sides; *praescutellum* not indicated; *scutellum* convex, narrowed to base, transverse anteriorly; *metanotum* very narrow, transverse, lower than mesonotum and epinotum; *epinotum* about as broad as long, narrowed slightly to base, dorsal surface convex, about as long as declivity, declivity slightly concave, angle