

Described from 24 workers (H.1426 A and H.1426 B) taken by Mr. H. E. Box in hollow stems (domatia) of mature *Canthium glabriflorum* inhabited by these ants.

Gold Coast, Bunsu, Jan. 1, 1945.

This distinct species comes in the *C. (A.) depressa* Latr. group, to which *C. (A.) mottazi* Santschi, from the Ivory Coast, also belongs.

*Crematogaster (Atopogyne) togoensis*, sp. n.

♀. Black, somewhat shining, clothed with short decumbent yellow hairs and some outstanding longer yellow hairs; club of antennæ and tarsi dirty yellow; mandibles reddish at apex.

*Head* a little longer than broad, broadest before posterior angles, which are round, posterior border excised in middle; *mandibles* triangular, striate and punctate, masticatory border armed with four rather strong teeth, the apical one longest and sharpest; *clypeus* convex, anterior border rounded, slightly sinuate at sides, longitudinally striate, posterior border rounded; *frontal area* narrow, triangular, well defined, but shallow, longitudinally striate; *frontal furrow* fine, continued beyond centre of head, spaces on each side very finely longitudinally striate, rest of head slightly more strongly striate and faintly and fairly closely punctured; *frontal carinæ* narrow, flat, divergent posteriorly; *antennal foveæ* fairly large and deep; *antennæ* 11-jointed, *club* 3-jointed, *scape* not reaching posterior border of head, not much thickened to apex, *funiculus* with first joint broader, but shorter, than second, club increasing in length and breadth, last joint about as long as the two preceding taken together. *Thorax* rather short, longer than broad, contracted at meso-epinotal furrow, sides rounded anteriorly to furrow; *pronotum* finely longitudinally striate and punctate, the punctures on the sides are larger than on the disc and wider apart, disc (with mesonotum) submarginate, humeral angles rather prominent; *mesonotum* rather flat, very finely longitudinally striate and punctured, the striæ on sides being stronger, a strong carina extends from the pointed anterior margin to the declivity at base, posterior border declivous and transversely striate; *meso-epinotal furrow* wide and deep; *epinotum* armed with two rather stout short sharp spines,