

projecting outwards, dorsal surface not as long as declivity, longitudinally striate, declivity somewhat concave, smooth and shining, transversely striate at sides; *post-petiole* broader than long, sides rounded, rather flat on disc, with a slight longitudinal impression posteriorly; *gaster* heart-shaped, first segment very finely and closely punctured, which gives a somewhat dull appearance, with larger (though quite small) punctures regularly scattered over it, second segment finely transversely striate and somewhat shining.

Long. 3-4 mm.

Type in B.M. (Nat. Hist.) Coll.

No. H.1385. Described from thirteen specimens taken by Mr. H. E. Box on *Cacao* (pod). French Togo, Mayondi (two miles east of Honuta, British Togo), Nov. 5, 1944.

Subfamily FORMICINÆ.

Camponotus (Myrmopiromis) flavosetosus, sp. n.

♂. Black, mandibles and legs brown, articulations of the legs, tarsi and funiculi lighter. Head and thorax dull, *gaster* shining, clothed with very sparse yellowish pubescence and fairly long outstanding pointed golden-yellow hairs.

Head triangular, narrowed to apex, slightly rounded to posterior angles, which are round, considerably broader behind eyes than anteriorly, posterior border almost straight, the sculpture consists of very fine, close, somewhat reticulate puncturation; *mandibles* strong, triangular, with larger and smaller scattered punctures, masticatory border armed with five or six strong teeth, the apical one being long and sharp; *clypeus* large, convex, triangular, broadly but distinctly carinate, anterior border widely excised in middle, rather deeply sinuate before side-pieces, furnished with a row of rather uneven yellow bristles or hairs, the centre one being the longest; *frontal area* indistinct; *frontal furrow* fine and narrow, extending beyond the centre of the flat surface between the frontal carinæ; *frontal carinæ* long, sharp, raised, sinuate in front and behind, flat surface between, broadest behind middle; *antennæ* 12-jointed, *scape* narrow, slightly curved, broadest before apex, reaching a little beyond posterior border of head, *funiculus* with all