

shallowly punctured, armed at apex with four teeth, the apical one long and pointed, the 4th blunt and bent backwards to form the 1st of a row of blunt teeth on masticatory border; *clypeus* with anterior border concave, finely transversely striate at sides, longitudinally so in centre which extends between the lobes of the front; *frontal carinæ* short, not very long, forming a scrobe at sides for the reception of part of the scape; *frontal furrow* rather deep, extending to median ocellus; *antennæ* 12-jointed, *scape* short, thick, sparingly and shallowly punctured, *funiculus* increasing in breadth to apex, joints 1-8 smooth and shining, very sparingly punctured, the last three joints, which form a somewhat flat club, rather closely punctured, especially the apical one which is bluntly pointed and is slightly longer than the other two joints taken together; *eyes* large, broad oval, rather prominent, situated on top of sides of head, behind the centre of the same; *ocelli* moderate. *Thorax* robust; *pronotum* with a neck, sides rounded, posterior border semicircular, faintly margined, embracing mesonotum, neck finely transversely striate, posterior portion of pronotum with larger and smaller scattered shallow punctures; *mesonotum* with a rather deep slanting furrow on each side forming a short triangle of anterior portion of disc, which is faintly punctured in front, *scutum* with large shallow scattered punctures, and faint longitudinal striæ, a narrow furrow extends from the point of triangle to præscutellum; *præscutellum* very narrow in centre; *scutellum* transverse, rather flat, with large, scattered, shallow punctures; *metanotum* narrow, slightly raised, with closer and larger punctures than scutellum; *epinotum* closely transversely striate, angle between dorsal surface and declivity not marked, declivity somewhat concave to base. *Petiole* transverse, sides rounded, anterior and posterior borders almost straight, with large shallow punctures, sparing on disc, more abundant at sides and front and hind borders; *post-petiole* transverse with a neck, which is finely transversely striate anteriorly, with short well-marked longitudinal striæ posteriorly, almost smooth on disc, with larger and smaller scattered punctures at sides, strangulation between post-petiole and gaster well marked; *gaster* almost smooth and sharply pointed to apex. *Legs* rather short, femora and tibiæ