

middle of sides of head; *ocelli* rather small; *antennæ* 12-jointed; *scape* short, curved, thickened towards apex, only extending slightly beyond posterior border of eye, *funiculus* with first joint slightly broader and longer than the following six joints, which are transverse; *club* 3-jointed, first joint shorter and narrower than the second, last joint longer than the two preceding taken together. *Thorax* narrowed in front and behind, broadest across humeral angles, flat on disc; *pronotum* transverse, narrowed to a neck in front, raised posteriorly, embracing the mesonotum, suture between pro- and mesonotum very distinct, anterior border of neck narrowly margined, raised, posterior portion of pronotum, smooth in middle, rather rugosely punctured at sides; *mesonotum* large, flat, narrowed anteriorly, puncturation as in head but somewhat finer, suture between scutellum and præscutellum very narrow but distinct; *præscutellum* only present at sides; *scutellum* transverse, flat, broad oval, anterior margin only separated from mesonotum by the fine suture, smooth in middle, puncturation as in mesonotum at sides; *metanotum* transverse, narrow, flat, slightly lower than the mesonotum, smooth in middle, with a row of five punctures at sides; *epinotum* about as long as broad, dorsal surface transverse, much shorter than declivity, smooth in middle, striate at sides, angle between dorsal surface and declivity very marked, declivity slightly concave, very smooth and shining, sides raised. *Petiole* longer than broad, rounded and smooth on upper surface, anterior surface flat and smooth, posterior surface sloping and slightly narrowed, beneath armed with a rather large blunt projection; *post-petiole* slightly longer than broad, upper surface rounded, with a few small scattered punctures, armed beneath with a large, blunt, tooth-like projection; *gaster* long, oval, rather flat, first segment very large and finely punctured. *Legs* moderate, *femora* clavate. *Wings* pale yellowish brown, *ptero stigma* and *veins* pale brown, one discoidal, a long cubital, and radial cell open, delicately fringed with short hairs.

*Long.* 4 mm.

Described from a single winged female, Nadzab, Markham R. val., New Guinea, 16th July, 1944. Taken by K. B. Krombein.