

Although this specimen is somewhat defective, the parts missing are not of much taxonomic importance in this genus, and as it is evidently a very distinct species, it appears best to describe it here.

♂. Brownish yellow, eyes black, apex of last four visible segments of gaster brown, body clothed with longer and shorter yellow outstanding hairs, apex of gaster and genitalia fringed with yellow hairs.

*Head* without eyes longer than broad, slightly narrowed behind, posterior angles bluntly rounded and margined, posterior border slightly excised in middle and finely margined; *mandibles* narrow and pointed, barely meeting when closed, apex armed with a thin, sharp tooth and a shorter, blunter one before it; *clypeus* convex on disc, and finely striate longitudinally, anterior border almost straight in middle and slightly sinuate at sides, finely margined, sides of disc also with raised fine margins; *frontal area* rather large, triangular, with two shallow hollows in front, not very distinctly defined, *frontal carinæ* slightly raised, with sharp narrow rims, slightly convergent and then divergent behind; *eyes* very large, round and prominent, situated a little in front of middle of sides of head; *ocelli* large, situated on vertex of head, which is somewhat raised; *antennæ* 12-jointed, scape and first joint of funiculus smooth and shining, the rest pubescent, *scape* about as long as the first three joints of funiculus taken together, first joint of *funiculus* shorter than second, last joint not quite as long as the two preceding taken together. *Thorax* longer than broad, narrowest in front and behind, rather robust; *pronotum* narrow, transverse, rather flat, lower than mesonotum, anterior angles very bluntly rounded; *suture* between pro- and mesonotum distinct; *mesonotum* convex, robust, longer than broad, narrowed in front and behind, broadest before insertion of forewing, anterior border rounded, posterior border almost straight; *mayrian furrows* very indistinct; *præscutellum* very narrow, transverse, flat; *scutellum* broad oval, transverse, convex but not very prominent; *metanotum* narrow, transverse, flat; *epinotum* convex, narrowed to base, angle between dorsal surface and declivity rounded, *dorsal surface* shorter than declivity, *declivity* convex rounded. *Petiole* longer than broad, longer than post-petiole, narrowed anteriorly, with a projection on