very slightly raised: eyes fairly large, round, not very convex, situated considerably in front of centre of sides of head; ocelli small, situated rather far back on vertex; antennæ 12-jointed, scape not nearly reaching posterior border of head, funiculus slightly thickened towards apex, 1st and 2nd joints subequal, longer than the following joints, the last is shorter than the one preceding. Thorax considerably longer than broad, broadest across insertion of forewings, narrowed considerably in front, less so behind; pronotum narrowed in front, widely and deeply excised posteriorly, embracing the mesonotum; suture between pro- and mesonotum very marked: mesonotum about as broad as long, broadest across middle, narrowed in front, posterior border straight; præscutellum flat, narrow, transverse: scutellum broad oval, not very convex nor projecting; mesonotum narrow, transverse, slightly raised, but lower than scutellum; epinotum rounded, convex, angle between dorsal surface and declivity not marked; dorsal surface longer than declivity. Petiole furnished with a rather low transverse node, rounded above; gaster rather long, and somewhat voluminous. Wings pale yellow, pterostigma and veins pale brown, one discoidal cell, one complete cubical cell, and one incomplete cubical cell, and rather long radial cell present.

Long. 5.7 mm.

Described from two winged females taken by K. V. Krombein, Nadzab, Markham R. val., New Guinea, July and August, 1944.

Iridomyrmex tigris Stitz. Gess. Naturf. Fr. 1912, p. 507. One winged \circ , Nadzab, Markham R. val., May, 1944.

VI Tribe ANONYCHOMYRMINI Tribe n. Male tarsi without claws.

√ ANONYCHOMYRMA, gen. nov.

Antennæ 13-jointed, scape short, maxillary palpi 5-jointed, labial palpi 3-jointed, no mayrian furrows present, the last joint of the tarsi not furnished with claws, but with hairs and the pulvilli large as in *Œco-phylla*.