Anonychomyrma myrmex, sp. n. VV

3. Black, legs brown, tarsi lighter, mandibles brownish yellow, whole body very finely punctured and pubescent,

gaster with short black outstanding hairs.

Head without eyes longer than broad, projecting posteriorly in a blunt wedge which supports the ocelli, posterior broder of which excised between lateral ocelli, sides behind eyes rounded, posterior border slightly excised; mandibles long, narrow, masticatory border armed with fine short teeth, the apical tooth sharply pointed, the tooth next to it longer and sharper than those following; clypeus transverse, convex, anterior border slightly rounded; posterior border bluntly pointed extending between insertion of antennæ; frontal area and frontal furrow slightly indicated; frontal carinæ short, sharp, slightly raised; eyes very large, prominent, round: ocelli very large and projecting; antennæ 13jointed, filiforme, all joints longer than broad, scape short, longer than first joint of funiculus, but shorter than second, first joint short. Thorax longer than broad, broadest at insertion of forewings, rather robust; pronotum very transverse, narrow, sides bluntly rounded at base and reaching the insertion of the anterior wings; mesonotum large, convex and prominent; præscutellum transverse, narrow, flat; scutellum slightly transverse, very convex and prominent: metanotum transverse, narrow, raised: epinotum large, round, prominent, angle between dorsal surface and declivity very slightly marked, declivity considerably shorter than dorsal surface. Petiole furnished with a round, narrow, slightly raised node, fitting close to gaster; gaster short oval. Legs fairly long, the last joint of the tarsi without clavus, but furnished with fine hairs and large pulvilli as in Œcophylla. Wings moderate, pale yellow, pterostigma and veins light brown; one discoidal cell, one large cubital and closed radial cell present, the cubitus not soldered to the radius, but their branches united by a short cubital nerve.

Genitalia: annula lamina transverse, narrow, almost straight at base, widely excised at apex; squamulæ rather long, somewhat narrow; stipites short, narrow, slightly curved, sharply pointed, hairy; volsellæ long and narrow, bent into a sharp hook at base; laciniæ curved