

and bluntly pointed; *sagittæ* thin, pointed to apex, finely serrate on inner margins.

Long. 4·2 mm.

I am indebted to Dr. van Emden for kindly making excellent preparation of the mouth-parts, tarsus, and genitalia for me.

Subfamily FORMICINÆ.

Tribe ECOPHYLLINI.

Ecophylla smaragdina F., subsp. *subnitida* Emery, Ann. Soc. Ent. France, lx. p. 565 (1891).

One winged ♀, Maffin Bay, Dutch New Guinea, October, 1944.

Tribe CAMPONOTINI.

Opisthopsis pictus Emery, var. *lepidus* Wheeler, Bull. Mus. Compar. Zool. lxii. p. 346 (1918).

Two ♀♀, Port Moresby, August 25th–28th, 1944.

Camponotus (Tanemyrmex) coxalis F. Smith, Journ. Proc. Linn. Soc. Lond. Zool. iii. p. 136 (1858).

One ♀ Nadzab, Markham R. val., June, 1944.

Camponotus (Colobopsis) desectus F. Smith, Journ. Proc. Linn. Soc. Lond. Zool. v. p. 97 (1861).

One winged ♀, Nadzab, Markham R. val., August, 1944.

Polyrhachis (Campomyrma) hecuba Forel, Rev. Suisse Zool. x. p. 527 (1902).

One winged ♀, Nadzab, Markham R. val., May, 1944.

Polyrhachis (Chariomyrma) aurea Mayr, var. *obtusus* Emery, Ann. Mus. Stor. Nat. Genova, xxxviii. p. 589 (1897).

One ♀, Nadzab, Markham R. val., June, 1944.

Polyrhachis (Chariomyrma) cingula, sp. n.

♀. Black, mandibles, antennæ, palpi, and legs red, teeth of mandibles and last four joints of tarsi blackish. In one specimen the upper surface of the spines and part of the explanate border of the pronotum are yellow. Whole body clothed with long outstanding yellow hairs.