

*Petiole* articulated to post-petiole for its whole breadth, anterior surface slightly concave, armed with a large blunt tooth beneath pointing downwards; *suture* between post-petiole and rest of gaster well marked; *gaster* oval, bluntly pointed at apex, not as long as thorax. *Sting* well developed. *Legs* moderate, *femora* compressed; posterior pairs of tibia armed with one spur, which is rather broad, broadest along apical border, which is finely pectinate.

*Long.* 2.7-3 mm.

♂. Shining blackish brown, legs and antennæ pale brownish yellow, first two or three of the joints of the funiculus paler; pubescence yellow, body clothed with short, scattered, outstanding brown hairs, less sparse on gaster.

*Head* somewhat globular, including eyes as broad as long, posterior angles rounded, posterior border almost straight; *mandibles* very narrow, pointed, meeting when closed; *clypeus* convex on disc, anterior border straight, narrowly margined, furnished with a row of very short hairs, posterior border pointed, and extending between the frontal carinæ; *frontal carinæ* much as in ♀; insertion of antennæ not covered; *eyes* very large, convex, slightly kidney-shaped; *ocelli* moderate, situated on vertex of head; *antennæ* 13-jointed, moderately long, very gradually increasing in length and breadth to apex, *scape* short, about as long as the two first joints of the funiculus taken together, first joint small, round, slightly broader, but shorter, than second joint, last joint longer than the two preceding joints taken together. *Thorax* narrowed in front and behind, broadest at insertion of forewings; *pronotum* transverse, furnished with a short neck; *mesonotum* convex in front, flat towards base; *mayrian* furrows present; *præscutellum* transverse, narrow; *scutellum* moderate, round oval, convex; *metanotum* narrow, transverse; *epinotum* long, angle between dorsal surface and declivity fairly distinct, declivity shorter than dorsal surface, slightly concave. *Petiole* pedunculate, articulated to post-petiole for its whole breadth, anterior surface flat, rounded above; *post-petiole* broader than petiole, suture between it and rest of gaster distinct, but not much constricted at sides; *gaster* long, oval, narrowed in front and behind; *epipygium* pointed.