

and *veins* pale yellowish brown; one large long *cubital cell*, a long closed *radial cell* present. The *cubitus* extends as far as the margins of the wing. The punctures of the *præscutellum* and *scutellum* are larger and more distinct than those of the rest of the body.

*Long.* 5.5 mm.

Described from a worker and a winged female. Maffin Bay, Dutch New Guinea, August, 1944. E. S. Ross Collection.

*Polyrhachis (Cyrtomyrma) dentata*, sp. n. (Fig. 3.)

♀. Black, shining, with the usual reticulations and very small punctures; palpi, trochanters, spurs, and claws brownish red; extreme tip of antennæ yellow.

*Head* broad, massive, broadest at base, narrowed anteriorly, broader at base than thorax, posterior angles rounded, posterior border excised and truncate in middle;

Fig. 3.



*Polyrhachis (Cyrtomyrma) dentata*, sp. n. Scale.

*mandibles* massive, masticatory border armed with four sharp teeth, the apical one being considerably the longest and sharpest; *clypeus* large, convex, anterior border truncate, and slightly raised, with a very small tooth at each corner; *frontal area* very indistinct; *frontal furrow* very fine and narrow, continued between the frontal carinæ almost to their base; *frontal carinæ* long, sinuate in middle; *eyes* large, round, not very convex, situated considerably behind middle of sides of head; no *ocelli* present. *Thorax* broad, massive, convex; *pronotum* furnished with a short transverse neck, slightly excised in front, and not very narrowly margined, shoulders rounded; *suture* between pro- and mesonotum fine and not deep, but very distinct; *suture* between meso- and epinotum not indicated; *epinotum* unarmed, the posterior border of the declivity deeply and narrowly margined; *scale* with four pointed teeth, the outer pair being long