

14. *Monomorium (Lampromyrmea) atomus* Forel subsp. *cospeis* Donisthorpe  
1 ♀, 3-5.V.35.

15. *Solenopsis cooperi* spec. nov.

♀. — Head, thorax and pedicel light brownish yellow, antennae, legs, and gaster pale yellow, eyes and teeth of mandibles black; whole smooth and shining, furnished with short scattered yellow outstanding hairs.

*Head* longer than broad, slightly broader anteriorly than posteriorly, posterior angles rounded, posterior border slightly and evenly excised, dorsal surface with some small scattered punctures, and a round shallow fovea or puncture on disc; *mandibles* powerful, faintly longitudinally striate, outer border rounded, inner border straight, masticatory border armed with 3 teeth, two sharp ones at apex, the apical one being the longer, and one blunt tooth at base; dorsal surface of *clypeus* convex, anterior border sinuate on each side, and armed in centre with two small blunt teeth; *frontal carinae* not long, narrow, raised, rather wide apart, and slightly divergent behind; *eyes* large, round, slightly convex, consisting of a number of facets, and situated in front of middle of sides of head; *ocelli* moderate; *antennae* 11-jointed, *scape* curved, not reaching posterior ocelli, *funiculus* with first joint longer than broad, longer than the following seven joints, which are transverse, last two joints large, elongate, forming a two-jointed club, the last joint of which is pointed and longer than the preceding. *Thorax* narrow, considerably longer than broad; *pronotum* transverse, short, furnished with a neck anteriorly; *mesonotum* higher than pronotum, which it slightly overlaps, longer than rest of thorax, narrowed and rounded anteriorly, *scutum* of mesonotum with a few small scattered punctures, slightly convex; *scutellum* round oval, rather flat on disc; *metanotum* narrow, transverse, lower than scutellum and epinotum; *epinotum* sloping, narrowed behind, no marked angle between dorsal surface and declivity. *Petiole* and *post-petiole* with dorsal surfaces subequal, convex, rounded above and at sides; *gaster* long, voluminous, narrowed in front and behind, truncate at base. *Legs* moderate. *Wings* transparent, veins and pterostigma pale yellow, one discoidal and one cubital cell present, radial cell open. *Long.* 4.8 mm.

♂. — Brown, smooth and shining, mandibles, antennae and legs pale yellow. Body furnished with short yellow outstanding hairs. *Head* small, triangular, posterior angles rounded, posterior border truncate; *mandibles* narrow, masticatory border armed with three teeth; *clypeus* convex; *frontal carinae* short, divergent, not close together; *frontal furrow* in the form of a rather deep small fovea; *eyes* kidney-shaped, very large and prominent, situated on front of sides of head; *ocelli* large, situated on top of posterior part of head; *antennae* 12-jointed, *scape* a little longer than first joint of funiculus,