der widely V-shaped, the furrow which separates it from the rest of the head rather deep; frontal carinæ forming the upper border of a rather deeply hollowed scrobe above the eyes; eyes large, round oval, ocelli small; antennæ six-jointed, fairly long; scape stout, curved, longer than the funiculus, very finely granulate, with a few adpressed short hairs on upper surface, anterior border with a row of longer and shorter not very close outstanding blunt spine-like hairs; funiculus covered with close adpressed silvery hairs pointing forwards, last two joints long and broader than those preceding, last joint pointed and longer than the two preceding taken together. Thorax stout, convex, very finely granulate; pronotum furnished with a neck anteriorly, transverse, posterior border semicircular, embracing the mesonotum; mesonotum large, prominent, about as long as broad, broadest just before base, posterior border almost straight, a very fine narrow suture separates it from præscutellum; præscutellum narrow, transverse, almost flat, separated from scutellum by a broad rounded suture which is deeper than that separating it from mesonotum; scutellum round oval, lozenge-shaped, transverse, somewhat prominent; metanotum very narrow, transverse, situated beneath scutellum; epinotum descending abruptly from beneath metanotum, armed with two short sharp teeth, space between the teeth somewhat hollowed out. Petiole with a long peduncle, and a node at apex rounded above and at sides, finely granulate; post-petiole finely granulate, broader than node of petiole, broadest at apex, rounded above and at sides. A white spongiform substance is present beneath the petiole and post-petiole and also along the junction between the petiole and postpetiole, and the post-petiole and gaster; gaster broadly oval, pointed at apex, broader in front than behind, broadest just before apex of first segment which is considerably longer than the rest of the gaster and very finely longitudinally striate. Rest of segments smooth and shining. Legs rather long, clothed with long decumbent glittering scale-like hairs. Long. 5 mm.

Redescription of F. Smith's type of Strumigenys man-