

Subfamily *PONERINÆ.*Tribe *PONERINI.*

*Ponera eduardi* Forel, Bull. Soc. Vaud. Sci. Nat., xxx, p. 15 (1894); Donisthorpe, Ann. Mag. Nat. Hist. (s. 12), iii (1950).

Known distribution: Basin of the Mediterranean, Madeira and Canaries.

6 ♀♀, West Turkey, Izmir, Denizli, No. 396, 3 ♀♀, Bebek, Istanbul, No. 131.

Tribe *PROCE R ATI INI.*

*Sysphincta europea* Forel, Ann. Soc. Ent. Belg., xxx, C.R. p. 162 (1884); Donisthorpe, Ann. Mag. Nat. Hist. (s. 12), iii (1950).

Winged and deälated ♀♀ and ♀♀, West Turkey, Bebek (C. Kosswig), 1947.

Subfamily *MYRMICINÆ.*Tribe *MYR M I C I N I.*

*Myrmica lobicornis* Nylander, Acta. Soc. Sci. Fennicæ, ii, p. 932 (1846).

Known distribution: North Europe, more in the south in the mountains, Alps, Appennines.

14 ♀♀, East Turkey, Nazit Gölü, February 2, 1947, Kars.

*Myrmica scabrinodis* Nylander, Acta. Soc. Sci. Fennicæ, ii, p. 930 (1846).

Known distribution: Europe, North and Central Asia.

Winged ♀, 4 ♀♀, Odemis, Izmir, No. 197; 1 winged ♀, Dinar Seyharap p'nari.

*Myrmica sulcinodis* Nylander, Acta. Soc. Sci. Fennicæ, ii, p. 934 (1846).

Known distribution: North of Europe and Asia, 4 ♀♀, Uludağ, 2200 m., Bursa, No. 19,