

*Messor barbarus* L., subsp. *semirufa* E. André, Spec. Hym. Europe, ii, p. 355 (1883).

Known distribution: Borders of the Caspian Sea, Syria, Persia, Abyssinia.

No. 213, common Antalya. No. 254, 3 ♀♀, W. Turkey, south coast of Anatolia. No. 255, 3 ♀♀, Araplarköy. No. 256, 6 ♀♀, south coast of Anatolia.

*Messor barbarus* L., subsp. *médiosanguineus* Donisthorpe, Proc. R. Ent. Soc. Lond. (B), xv, Pts. 5 and 6, p. 53, (June 1946).

Very many ♀♀, Gaziantep, August 25, 1947. The subspecies was described on 21 specimens 2 1/2 and ♀♀ taken in Turkey previous to 1946 by Professor C. Kosswig.

*Messor barbarus* L., var. *meridionalis* E. André, Spec. Hym. Europe, ii, p. 355 (1883).

Known distribution: Greece, Albania, Istanbul, Algeria, Tunis.

No. 86, 1 ♀, Bebek, Istanbul. No. 162, Toros, 13 ♀♀. No. 125, 5 ♀♀, Elbistan. No. 226, 26 ♀♀ and 2 1/2, Ankara.

*Messor oertzeni* Forel, var. *amphigea* Forel, Bull. Soc. Vaud. Sci. Nat., xlvi, p. 347 (1911).

No. 388, 1 ♀, West Turkey, Sarayköy, November, 1945 (C. Kosswig). No. 230, 11 ♀♀, Ankara, Istanbul. No. 213, ♀♀ and 2 1/2, Antalya.

*Gonioma burri* Donisthorpe, Ent. Record, lxiii, p. 60 (1950).

One winged ♀, two deälated ♀♀, West Turkey, Black Sea coast, Kopa.

*Ceratopheidole pergandei* Donisthorpe, Ann. Mag. Nat. Hist. (s. 12), iii, p. 639 (1950).

5 ♀♀, S.E. Turkey, Gaziantep, August 25, 1947.

#### Tribe CARDIOCONDYLINI.

No. 34, *Cardiocondyla elegans* Emery, Ann. Accad. Aspir. Nat. Napoli, ii, p. 21 (1869).

Known distribution: Italy, Central France, Spain, Caucasus. 3 ♀♀, Ankara, Istanbul.