

## Tribe C R E M A T O G A S T R I N I .

*Crematogaster (Orthocrema) sordidula* Nylander, Ann. Soc. Sci. Fennicæ, iii, p. 44 (1849).

Known distribution : Basin of the Mediterranean and Central Asia. No. 44, 5 ♀♀, Armutlu, east coast of Marmara.

*Crematogaster (Acrocaelia) scutellaris* Olivier, Encycl. Méthod. Insect. vi, p. 497 (1791).

Known distribution : Central Europe and east of the Adriatic, Algeria, Tunis, Caucasus.

No. 42, 7 ♂♂, 10 ♀♀, Baltalimani, Istanbul. No. 138, 24 ♀♀, Bebek, Istanbul.

## Tribe S O L E N O P S I D I N I .

*Solenopsis fugax* Latreille, Fourn. Fr., p. 46 (1798).

Known distribution : Southern and Central Europe, England, Sweden, Morocco, Asia, West and Central Japan.

No. 63, over 20 ♀♀, Kara, Istanbul. 5 ♀♀, Dinar, Layharah pinari. No. 165, 1 ♂, 1 winged ♀, and 1 deälated ♀, Yeşilkoy, Istanbul.

*Monomorium (Holcomyrmex) glabrum* E. André, Spec. Hym. Europe, ii, p. 345 (1883).

Known distribution : All India, Ceylon, Burma.

10 ♀♀, West Turkey, Vilayet, Bardur, Gulda Gölü, October, 1946.

## Tribe P H E I D O L I N I .

*Pheidole (Pheidole) megacephala* Fabricius, Ent. Syst., ii, p. 361 (1793).

Cosmopolitan. In hothouses.

No. 370, West Turkey, Izmir, Denizli, 5 ♀♀. No. 256, 6 ♀♀, Araplarköyü, south coast of Anatolia.

*Pheidole (Pheidole) pallidula* Nylander, Act. Soc. Sci. Fennicæ, iii, p. 42 (1849).

Known distribution : West Basin of the Mediterranean in Europe and in Africa, limited in Europe by the Adriatic.