

No. 443, 5 ♀♀, many ♀♀, Ankara, August 2, 1945.
 No. 285, West Turkey, Cihanbeyli, Istanbul, 23 ♀♀.
 No. 222, 4 ♀♀, 4 ♀♀, Odemis, Izmir. No. 3, 1 ♀♀, 4 ♀♀,
 Ankara, Bursa. No. 78, 11 ♂♂, 6 ♀♀, 30 ♀♀, Bebek,
 Istanbul. No. 163, 18 ♀♀, Yeşilköy, Istanbul. No. 211,
 1 ♀♀, 7 ♀♀, Antalya, Izmir. No. 203, 2 ♀♀, 7 ♀♀, ditto,
 ditto.

Tribe LEPTOTHORACINI.

Leptothorax bulgaricus Forel, subsp. *græca* Forel, Bull.
 Soc. Vaud. Sci. Nat., xlvii, p. 336 (1911).

Known distribution : Greece, Dalmatia.

No. 190, 5 ♀♀, Armutlu, east coast of Sea of Marmara.

Tribe TETRAMORINI.

Tetramorium cæspitum Linnæus, Syst. Nat. (ed. 10), i,
 p. 581 (1758).

Known distribution : The whole of Europe and north
 of Asia, Japan, imported into Africa and North America.

Common and widely distributed. We have specimens
 from Edremit, Adilcevaz, Isikli, Kars, Kopa, Uludağ,
 etc. Nos. 2, 9, 11, 20, 53, 61, 71, 220.

Tetramorium cæspitum L. var. *brevicornis* Emery, Bull. Soc.
 Ent. Ital., xlvii, p. 194 (1916).

Known distribution : Sardinia, Corsica.

No. 947, 1 ♀, Turkey, Black Sea coast, Zondulkak, May,
 1947

Tetramorium cæspitum L. subsp. *semilævis* E. André,
 Spec. Hym. Europe, ii, p. 286 (1882).

Known distribution : Basin of the Mediterranean,
 Central Asia.

9 ♀♀, very dark, but not quite as black as the var. *niger*,
 South-east Turkey, Gaziantep, August 25, 1947. 50 ♀♀,
 West Turkey, Aigol Gomic, October, 1946. No. 20, 8 ♀♀,
 Uludağ, 1900 m.

Subfamily DOLICHODERINÆ.

Tribe TAPINOMINI.

Tapinoma erraticum Latreille, Ess. Fourmis France, p. 24
 (1798).

Known distribution : Europe, middle and southern ;
 shores of the Mediterranean, Caucasus, Central Asia.