

10 ♀♀, Gaziantep, Syrian frontier, August 25, 1947, 2 ♀♀, Ahlat, June 30, 1947. No. 139, 17 ♀♀, West Turkey, Armutlu, east coast of Sea of Marmara. No. 39, many ♀♀, ditto, ditto. No. 31, 10 ♀♀, Ankara. No. 212, 12 ♀♀, Antalya, south coast of Anatolia.

Subfamily *FORMICINÆ*.

Tribe *PLAGIOLEPIDINI*.

*Plagirolepis pygmæa* Latreille, Ess. Fourmis, p. 45 (1798).

Known distribution: Temperate Europe, Italy and Italian Isles as far as the Balkans.

13 ♀♀, Gaziantep, Syrian frontier, August 12, 1947. No. 43, many ♀♀, Baltalimani, Istanbul. No. 128, 7 ♀♀, Nurhakköyü, Elbistan.

*Acantholepis frauenfeldi* Mayr, Verh. Zool. Bot. Ver. Wien, v. p. 378 (1855).

Known distribution: Dalmatia, etc.

7 ♀♀, coastlands Sogukoluk, February, 1946. No. 237, 6 ♀♀, Armutlu, east coast of Sea of Marmara. No. 275, 5 ♀♀, Mersin, south coast of Anatolia. No. 38, 35 ♀♀, Armutlu. No. 225, 1 deälated ♀, 60 ♀♀, Yeşilköy, Istanbul. No. 87, a deälated ♀, 40 ♀♀, Bebek, Istanbul. No. 164, 9 ♀♀, Yeşilköy, Istanbul.

Tribe *ACANTHOMYOPSIINI*.

*Acanthomyops (Donisthorpea) niger* Linnæus, Syst. Nat. (ed. 10), i, p. 580 (1758).

Known distribution: Europe, Caucasus, Siberia, Japan. 60 ♀♀, Gaziantep, Syrian frontier.

*Acanthomyops (Donisthorpea) alienus* Forster, Hym. Stud. i, pp. 36, 71 (1850).

Known distribution: Europe, Central and Southern Asia, Japan. Common and widely distributed.

We have specimens from Ankara, Armutlu, Baltalimani, Bebek, Hopa, Kars and Uludağ. Nos. 8, 10, 13, 18, 67, 48, 42, 85, 219 and 229.

*Acanthomyops (Donisthorpea) alienus* Forster, var. *alieno-niger* Forel, Fourmis Suisse, pp. 47, 49 (1875).

Known distribution: Europe.

No. 371, 2 ♀♀, Isikli, West Anatolia.