

No. 14, 4 ♀♀, Uludağ, 1900–2000 m. 5 ♀♀, East Turkey Nayik Gölü, July 2, 1947.

*Formica rufibarbis* Fabricius, Ent. Syst. ii, p. 355 (1793).

Known distribution : Europe, north of Asia, Himalayas.

11 ♀♀, 1 ♂, South-east Turkey, Gaziantep, August 25, 1947. No. 318, 1 ♀, Konya, coast of Anatolia. No. 12, 1 dealated ♀, Uludağ, Bursa, 1900 m. No. 32, 1 winged ♀, and 1 ♀, Ankara, Istanbul. No. 15, 5 ♀♀, Uludağ, 1700 m. No. 7, 3 ♀♀, Uludağ, 1900 m. No. 229, 4 ♀♀ Ankara, Istanbul.

*Formica fusca* Linnæus, Syst. Nat. (ed. 10), i, p. 580 (1758).

Known distribution : Europe and Asia, south and middle, in the Alps to 2400 m.

No. 51, 2 ♀♀, Uludağ, 1900 m. No. 8, 1 ♀ Uludag, 1900 m. No. 76, 1 large ♀, 5 small ♀♀, Uludağ, 1800 m.

*Formica fusca* L. var. *rubescens* Forel, Ann. Soc. Ent. Belg. xlviii, p. 423 (1904).

Known distribution : Europe, middle ; Italy.

No. 217, 26 ♀♀, Hopa, Haradeniz.

*Cataglyphis albicans* Roger subsp. *livida* E. André, Ann. Soc. Ent. France (6) i, p. 58 (1881).

Known distribution : Syria.

7 ♀♀, East Turkey, Ahlat, June 30, 1947.

*Cataglyphis cursor* Fonscolombe, Ann. Soc. Ent. France, (2), iv, p. 41, Bull. 69 (1846).

Known distribution : south of France.

No. 38, 2 ♀♀, Armutlu, east coast of Sea of Marmara. No. 221, 6 ♀♀, Odemis, Izmir. No. 37, 8 ♀♀, Ankara, Istanbul. No. 54, 30 ♀♀, Kars on the Causasian frontier.

*Cataglyphis bicolor* Fabricius, Ent. Syst. ii, p. 351 (1793).

19 ♀♀, South-east Turkey, Gaziantep, August 25, 1947. 7 ♀♀, East Turkey, Ahlat, June 30, 1947.

*Cataglyphis bicolor* F. var. *nodus* Brullé Exped. Sci. Morée Zool. ii, p. 326 (1832) (= *Myrmecocystus viaticus* Emery and Forel, var. *orientalis* Forel, and *M. viaticus desertorum* Emery, var. *orientalis* Emery).

11 ♀♀, South-east Turkey and Gaziantep, August 25, 1947.