

Cataglyphis bicolor F. var. *laevior* Stitz, Mitt. Zool. Mus. Berlin, viii, p. 348 (1916).

Known distribution : Algeria.

16 ♀♀, West Turkey, Vilazet Burdar, Gulda Gölü, October 1946.

Cataglyphis bicolor F. subsp. *megalocola* Förster, Verh. Naturh. Ver. Preuss, Rheinl. vii, p. 790 (1850).

Known distribution : Algeria.

No. 160, 1 small and 2 large ♀♀, Bürücek near Boganti, Toros.

Tribe CAMPONOTINI.

Camponotus (*Camponotus*) *herculeanus* Linnæus, Syst. Nat. (ed. 10) i, p. 599 (1758).

Known distribution : Europe, north and middle (mountains).

No. 47, 1 deälated ♀, Uludağ, 2000 m.

Camponotus (*Camponotus*) *higniperda* Latreille, Fourmis, p. 88 (1802).

Known distribution : Europe, north and middle, in the mountains in South Europe.

No. 16, 9 ♀♀, Uludağ, 1600 m.

Camponotus (*Camponotus*) *vagus* Scopoli, Ent. Carniol, p. 312 (1763).

Known distribution : Europe, south, Isles of Mediterranean.

Nos. 8 and 11, 28 ♀♀, Uludağ, Bursa. No. 949, 4 ♀♀, Black Sea coast, Zondulkak, May 1947. 11 ♀♀, East Turkey, Tetvan Nemizut, June 31, 1947. No. 79, 17 ♀♀, Büyükada, Principio, Isle of Princes. No. 325, 7 ♀♀, Konya, coast of Anatolia.

Camponotus (*Tanæmyrmex*) *æthiops* Latreille, Ess. Fourmis France, p. 35 (1798).

Known distribution : Europe, south and parts of temperate central, Caucasus.

No. 4, 1 ♂, 2 ♀♀, Uludağ, Bursa.