

versity; no. (2), British Museum; no. 1700a, University of Colorado; no. 2, Wickham coll.; no. 22,973, A. M. N. H.

The holotype specimen is a dorsal view of the insect with the wings spread. The ocelli, although probably present in the species, are not visible because of the sculpturing on the head. The venation of the forewing is variable with respect to the shape of the discoidal cell and the point of divergence of the first intercubitus.

The worker of this species is represented, I believe, by two specimens of fair preservation. One of these is a lateral view of the thorax and gaster, and a dorsal view of the head; the other (ergatotype) is entirely a lateral specimen. No sculpturing is discernible but I infer that this is due to insufficient preservation. The head, thorax, and petiole are identical with those of the above female. The measurements of the ergatotype are as follows: length of specimen, 6.0 mm.; length of head, 1.3 mm.; thorax, 1.7 mm.; gaster, 3.0 mm. Width of head, 0.9 mm.

*Ergatotype*.—No. 2803, M. C. Z. (S. H. Scudder).

*Paratype*.—No. 10,002, Peabody Museum.

#### DOLICHODERUS ROHWERI, sp. nov.

Plate 4, fig. 1. Plate 9, fig. 7. Plate 10, fig. 3

*Female*.—Length, 5.3 mm. Moderately slender; head somewhat longer than broad, posterior margin only slightly concaved, posterior angles rounded; lateral margins curved; clypeus probably quite large; scape short, not reaching the posterior margin of the head; funicular segments 2-10 about as long as broad, the first and last somewhat longer; eyes oval, of moderate size, situated at the middle of the sides of the head; thorax about as broad as the head; epinotum only slightly concaved posteriorly, and not extending backward far enough to project over the petiole; scale of the petiole with its anterior face nearly vertical and the posterior face at about a 45-degree angle; gaster about two and one-half times the length of the head. Clypeus striated, the entire head and pronotum finely reticulate, the sculpturing on the epinotum coarser; posterior face of the scale of the petiole with a few longitudinal striations. Length of head, 1.2 mm.; scape, 0.6 mm.; funiculus, 0.9 mm.; thorax, 1.7 mm.; gaster, 2.4 mm. Width of head, 0.85 mm.

*Holotype*.—No. 2801, M. C. Z. (S. H. Scudder).

*Paratypes*.—No. 10,001, Peabody Museum; no. 2825, Princeton