

9. *Ænictus* (*Ænictus*) sp.

Three colonies taken from the Cuernos de Negros Mountains at 1,500, 2,500, and 3,600 feet altitude. These are all related to the *turneri* Forel group in color, but differ in size and sculpture. I am unable to place them at this time.

10. *Ænictus* (*Typhlatta*) *alticola* Wheeler and Chapman.

Polis Pass, Benguet Province, at 6,000 feet.

11. *Ænictus* (*typhlatta*) *laeviceps* Fr. Smith.

Mt. Maquiling, Leyte, Cuernos de Negros, and several places on Negros Island, and on the island of Mindanao. In some colonies the workers seem to be more robust than in other colonies.

12. *Ænictus* (*typhlatta*) *luzoni* Wheeler and Chapman.

Bagia, Luzon, (Banks) 1913; Mt. Maquiling, Cuernos de Negros at 1,500 feet, and the Lake region north of the Cuernos at 2,700 feet.

13. *Ænictus* (*typhlatta*) *martini* Forel.

Mt. Maquiling: San Jose, Mindoro; Cuernos de Negros; throughout the island of Mindanao, (Gingoog, Davao, Lanao, Parker Lake on south Mindanao, and Zamboanga Peninsula). This species is more abundant than *laeviceps* Smith. The ratio in colonies on the Cuernos is 1 *laeviceps* to 4 *martini*.

*Martini* has small workers, which are abundant enough to attract attention to any one watching a foray. Measurements of the regular *martini* range from 3.4 to 3.55 mm. The range of ten small workers was 2.7 to 2.96 mm.

*Martini* Forel is the only species of *Ænictus* of which all three castes have been taken together.

LIST OF *ÆNICTUS* MALES TAKEN AT LIGHTS

1. Banks, Philippines, 1913, 1 male, no locality given.
2. Cuernos de Negros, 1922, 1 male at 1,500 feet.
3. Cuernos de Negros at 1,500 feet, 3 males at 1,500 feet.
4. Cuernos de Negros, male in pitcher plant at 4,000 feet.
5. Cuernos de Negros, 1925, 1 male at 1,500 feet.
6. Cuernos de Negros, April, 1934, 1 male at 1,500 feet.
7. Dumaguete City, 1934, 1 male.
8. Dumaguete City, October 1947, 1 male.
9. Iloilo City, 2 males taken.
10. Los Baños, Williams, 2 males.
11. Mt. Maquiling, Baker, 1 male.
12. Mt. Apo, Mindanao, Clagg, 6,000 feet, 1 male.
13. Limay, Luzon, 1 male.
14. Samar, McGregor, 1 male.
15. San Jose, Mindoro, Ross, 1 male.
16. Iligan, Mindanao, Baker, 1 male, described by Memozzii, as *Ænictus bakeri* Menn. (1925).