2. Antennal scapes with erect hairs; (body black or very dark brown, thorax sometimes reddish)neogagates lasioides var. vetula Wheeler
Antennal scapes without erect hairsneogagates Emery
3. Anterior clypeal margin notched in the middle
4. Gaster strongly shining, with very short sparse pubescence 5 Gaster opaque or subopaque, with longer dense pubescence 6
5. Erect hairs on head, thorax and gaster long and dense; clypeal notch indistinct in small workersperpilosa Wheeler
Erect hairs on head, thorax and gaster shorter and sparser; clypeal not distinct in small workers
6. Hairs on dorsal parts of body abundant, conspicuous, glistening white, obtuse or clavateobtusopilosa Emery
Hairs on dorsal parts of body sparse and more slender
7. Hairs nearly always absent from thoracic dorsum and petiolar bordersanguinea subnuda Emery
Hairs present though sparse on thoracic dorsum and petiolar border
8. Posterior corners of head shining; head, thorax and appendages yellowish red, gaster brownsanguinea puberula Emery
Posterior corners of head opaque or subopaque; thorax, petiole, appendages and lower surface of head dull, yellowish brown, vertex and occiput piceous brown, gaster deep piceous brown
9. Body robust; head of largest workers not or scarcely longer than broad; funicular segments 2-3 longer and more slender than segments 6-8; petiole usually with a rather sharp superior border; (body opaque; color light or dark red, with brown or black gaster)
10. Female larger than largest workers, measuring 6-11 mm.
Female not larger and sometimes even smaller than largest workers, measuring only 4-6 mm. microgyna var. rasilis Wheeler
11. Antennal scapes with erect hairs 12 Antennal scapes without erect hairs 13
12. Head and thorax bright yellowish red, legs reddish brownoreas Wheeler Red parts of body darker, legs dark brownoreas var. comptula Wheeler
13. Frontal area opaque
14. Erect hairs absent from gula and upper surface of head and thorax
15. Small forms (4-6.5 mm.)
16. Gaster blackish brown
17. Erect hairs on middle and hind tibiae confined to a double row of b-istles along flexor surface
Erect hairs on middle and hind tibiae usually abundant and covering all sur- faces, but at least there is a number of erect hairs besides the bristles on the flexor surface
18. Gaster densely clothed with short erect hairs which form a thick investiture mucescens Wheeler
Erect hairs on gaster widely spaced, not forming a thick investiture
rufa haemorrhoidalis Emery