

na *: Cassá de la Selva, Collsuspina (F); La Junquera (Y). Huesca *: Arguis, Torla (Y). Madrid: El Pardo (F), Sierra de Guadarrama (Y), Torrelodones (F). Málaga: Ronda (Y). Pontevedra *: Bayona, La Guardia, Ría de Vigo (F). Santander *: Potes (F). Segovia: Marugán (Y). Tarragona: Montblanch (F). Teruel *: Albarracín (Y). Zamora *: Zamora (Y).

Portugal.—Estremadura: Buçaco (W).

108. **Camponotus (Myrmosericus) micans** (Nyl.).

Huelva: Gibraleón (Y). Madrid: El Pardo (F), Montarco (M). Málaga: Ronda (Y). Sevilla: Los Palacios (Y).

Portugal.—Alto Alemtejo: Elvas (Y). Baixo Alemtejo: Beja (Y). Palma: Setubal (L).

This is widely distributed in South Spain. It is evidently listed by Ceballos as *C. rufoglaucus* Jerdon.

109. **Camponotus (Tanaemyrmex) aethiops** (Latreille).

Barcelona: San Feliú (B). Gerona: Collsuspina, Gerona (F). Huesca *: Jaca (C), Torla (Y). La Coruña: Corrubedo (F). Logroño: Monasterio de Valvanera en Anguiano (F). Lugo *: Sierra de Ancares (F). Madrid: El Escorial (G), Sierra de Guadarrama (F). Murcia *: Puerto Lumbreras (Y). Navarra *: Roncal (Y). Oviedo *: Puente de los Fierros (F). Pontevedra *: Isla Cies del Norte, Isla Cies del Sur, Isla de Ons, Pontevedra, Ría de Vigo (F). Santander *: Santoña (F). Soria *: Cabrejas del Pinar (Y). Tarragona *: Sierra de Montsant (F). Teruel *: Aliaga (F). Zaragoza: Embalse de Yesa (F).

110. **Camponotus (Tanaemyrmex) marginata** (Latr.).

Madrid *: El Escorial (M).

111. **Camponotus (Tanaemyrmex) pilicornis** (Roger).

Barcelona: San Feliú (B). León *: Sierra de la Cabrera (F). Madrid: Sierra de Guadarrama (F) (A) (C). Pontevedra *: La Guardia (F). Segovia *: Marugán (Y). Tarragona: Sierra de Montsant (F). Zaragoza: Sierra de Alcubierre (F).

Portugal.—Douro: Villa Nova de Gaia (W). Estremadura: Buçaco (W).

This species was also recorded by Roger (1863) from the Balearic Islands but this old record was not listed by Bernard (1956).