

*Camponotus (Myrmophyma) ephippium*, Sm., s.sp. *narses*, For.

♂. (Not described.) Length 11 mm. (♀ 6-8.5 mm.).

Black; head, except a brown triangular patch, with its apex at the ocelli, deep red; mandibles darker red, legs brown, the posterior femora sometimes red. Borders of gastric segments testaceous.

The grey pilosity and pubescence are similar to that of the ♀, but the pilosity is shorter. Tibiæ and scapes without erect hairs.

Head very thick and massive, broader than the thorax, the vertex much swollen; the head is as broad as long, broadest at occiput, where the angles are only slightly rounded, almost square, and prominent in profile; the sides almost straight as far as the lower cheeks, where they converge; occipital border widely and shallowly concave. There is a slight depression between the bulge of the cheeks and the frontal carinæ; the latter are nearly twice as wide apart behind as in front. Mandibles thick, with six teeth. Frontal groove distinct, reaching to a pit representing the anterior ocellus. Clypeus flatter than in the ♀, feebly carinate, the anterior border slightly concave in the centre, and crenulate as in the ♀. The scapes extend beyond the occiput by as much as their width.

Pronotum very broad, twice as broad as long, the mesonotum considerably narrower, convex in both directions; meso-epinotal suture distinct but faint, the base concave as in the ♀, but shorter in proportion; the declivity half as long as base, more vertical than in the ♀, and slightly concave.

Petiole in the form of a scale, not a node as in the ♀, broad, bluntly angular at top; in profile the top is bluntly pointed. Gaster nearly as broad as head.

Moderately shining. Mandibles coarsely punctate-striate. Whole of upper surface of head covered with somewhat elongate punctures, with a fine dense ground reticulation. The sculpture becomes feebler and the punctures more scattered at the sides and at the occiput, where the surface is shining. Pro- and mesonotum with a fine ground reticulation and a few punctures, epinotum more densely reticulate and matt. Gaster more abundantly provided with small punctures and microscopically reticulate. Legs with scattered punctures.

Armadale, W. Australia (*Clark*, nos. 222, 225, 268, 269).

*Camponotus (Dinomyrmex) RUFUS*, sp. n.

♂. Length 15-17 mm.

Rufous, gaster black. Head darker red than the thorax, and the legs more inclined to yellow than the thorax. Mandibles and clypeus somewhat fuscous. Ocellar region fuscous.

Body with a few short stiff hairs. Scapes without erect hairs, tibiæ with a few bristles on their under surface. On the mandibles and anterior border of clypeus are short golden hairs.

Head broader than long (5.2 by 4.4 mm.), broadest at level of eyes, the sides slightly convex, the occipital border widely concave,