

than the front face, hence, in profile the node presents a wedge-shaped rather than an arcuate outline.

The distributional records for *apache* are presented below. Unless otherwise noted all specimens were taken by the writer. The elevations of the majority of these records were checked with topographic sheets and an altimeter, the remainder with one or the other.

ARIZONA:

Huachuca Mountains: Ramsey Canyon (5500') single deälated female; Garden Canyon (5800') three colonies in *Quercus emoryi*; Carr Canyon (5400') two colonies in *Q. emoryi*.

Santa Rita Mountains: Madera Canyon (5900') one colony in *Q. emoryi*; Sweetwater (5800') three colonies in *Q. emoryi*.

Dragoon Mountains: Cochise Stronghold (5200') one colony in *Q. emoryi*.

Chiricahua Mountains: Chiricahua National Monument (5400') one colony in *Q. emoryi*.

Peloncillo Mountains: Cottonwood Canyon (4800') two colonies in *Q. emoryi*.

Baboquivari Mountains: Baboquivari Canyon (3500') one colony in *Q. oblongifolia*; Brown Canyon (4200-4400') type locality, nine colonies in *Q. emoryi* or *Q. oblongifolia*.

Ajo Mountains: (Organpipe Cactus National Monument) Alamo Canyon (2200-2800') ten colonies, eight in *Prosopis juliflora*, two in *Q. turbinella*.

Growler Mountains: (Organpipe Cactus National Monument) La Abra Wash (1300') two colonies in *Prosopis juliflora*.

Canelo Hills: Canelo Pass (5300') one colony in *Q. emoryi*.

Pena Blanca Springs (3900') Santa Cruz Co. Coll. L. F. Byars, one colony without data on nest site.

San Miguel (2400') Pima Co. Coll. L. F. Byars, one colony in *Prosopis juliflora*.

Organpipe Cactus National Monument Headquarters (1650') Coll. E. R. Tinkham, one colony, said to have been taken emerging from a rodent's burrow!