

rectangular, above with a faint median longitudinal elevation, its pilosity reclining and regularly arranged. Gaster very slender, the intersegmental areas bulging, very smooth and shiny. The rest of the integument is transversely chagreened and is covered with dense, short and adpressed hairs; on the anterior edge of each intersegmental membrane there is a row of longer hairs, reminiscent of the pilosity of *E. esenbecki*. Anal plate with four distinct points, the inner pair shorter than the outer. Claws simple as in the genus *Neivamyrmex*. — Holotype in the authors possession. Type locality: the border of Bolivia and Peru, from an unspecified locality. Other specimens have been found in the High Andes in the Satipo Area. Mr. Felix Woitkowsky coll."

Neivamyrmex detectus Borgmeier ♂

(Taf. 38 Fig. 4; Taf. 42 Fig. 16; Taf. 47 Fig. 11-12, 23; Taf. 49 Fig. 25; Taf. 70 Fig. 4)

Neivamyrmex detectus Borgmeier, 1953 p. 25, fig. 7, 15 (Maennchen; Itatiaia, etc.).

Typen: Maennchen (Holotype u. 19 Paratypen; CTB). Typenlokalitaet Itatiaia, Staat Rio de Janeiro, Brasil, J. F. Zikán leg.

Maennchen. — Zur Originalbeschreibung füge ich folgende Messungen einer Paratype vom Itatiaia hinzu. Kopfbreite 2,12 mm, Kopfhöhe 1,2 mm, Stirnbreite 1,08 mm. Augenlänge 1,04 mm. Breite des vorderen Ocellus 0,32 mm, Abstand der hinteren unter sich 0,52 mm, vom vorderen 0,16 mm, vom Auge 0,08 mm. Mandibeln 0,88 mm. Scapus 0,8 mm, Geissel 4 mm. Thorax: Laenge 4,4 mm, Breite 2,2 mm. Petiolus: Breite 1,4 mm, Laenge 1,4 mm. Gaster 8-8,5 mm. Hinterschenkel 2 mm, Hintertibiae 1,8 mm. Klauen fein gezähnt. Fluegel 13 mm, Breite 5 mm.

Arbeiter und Weibchen unbekannt.

Geographische Verbreitung. — Brasilien: Staat Rio de Janeiro und São Paulo.

Untersuchtes Material. — Holotype und 19 Paratypen, wie folgt: Rio de Janeiro, Itatiaia [Zikán 6.XII.28, 11.XII.28, 3.XI.29, 9.XI.31, 9.XI.37, XI.49]; Jussaral [Lopes & Lent 26.X.35]; Angra dos Reis [Travassos, Oiticica & Lins IX.36] [X.38]; Petrópolis [Borgmeier 3.XI.48]. São Paulo: Alto da Serra [Spitz IV.29, VI.29]; Ipiranga [Reh]; Cantareira [Luederwaldt IV.20; Menozzi det. *gravenhorsti*]; Eugênio Lefèvre [Travassos, Lopes & Oiticica 1.XI.37].