

reddish-brown. Thoracic sutures and posterior edges of the abdominal segments not darkened. Head without the mandibles nearly twice as broad as long; eyes very large and convex, occupying the entire sides of the head; with sinusoid inner borders. Mandibles relatively short, shorter than the space between their bases; thick in the basal  $1/2$ , tapering in the apical  $1/2$ ; smooth and shiny, with copious long hairs on the inner and outer borders. Scapes extending to the middle of the median ocellus, stout, thickest in the middle. First funicular joint as long as broad; all joints covered with dense, short, semierect hairs. Frontal carinae far apart, ending in front of the median ocellus in rounded elevations. The sharp carinae on the cheeks dark brown; in front with triangular foveae. The entire head smooth, with very few piligerous foveae on the vertex; shiny. Ocelli large, nearly touching the eyes (separated from the eyes by  $1/10$  their diameter). Prothorax very gibbous, the head barely visible from above, covered with dense, long and adpressed hairs. These hairs are much shorter than in the other species of *Woitkowskia*. Thorax very shiny. Petiole broader than long and nearly flat above, excavated anteriorly and above; its lateral margins well marked but rounded; completely covered with short and adpressed hairs, except for the excavated part, which has long and erect hairs, and the posterior edge of the petiole, which is densely covered with long and backward directed hairs, hiding its general outline. Gaster very elongate, shiny, with small piligerous foveae. Subgenital plate short and stout with curved inner points. — Holotype in the authors possession. Type locality: Vienna, Peru. Mr. F. Woitkowski coll.”.

*Woitkowskia bureni* Enzmann (1952). — “Male. Close to the species *W. kuertii* but larger. Length 15 mm. Color darker. The wings are brownish and have very dark brown veins. (In *W. kuertii* the veins are whitish). Head shaped as in *W. kuertii* but the lighter colored area on it is larger, leaving only the occiput dark colored. The entire head covered with abundant hairs (in *W. kuertii* only the vertex with abundant hairs). Other characters as in *W. kuertii* except for the sculpture on the gaster. *W. bureni* has the segmental borders densely and finely punctate while in *W. kuertii* they are smooth and shiny. — Holotype in the authors possession. Type locality: Satipo, Peru. Mr. Woitkowski coll.”.