

covered with pale erect hairs, which are somewhat longer than the distances between them. Head inclined ventrally more or less at a right angle to the body. Mouth parts feebly chitinized and pale yellowish. Mandibles subtriangular, very long and sharp-pointed, with the apex slightly curved. Maxillae with a lateral conical papilla, which is rather large and slightly curved. Differing somewhat from the common ponerine type and showing a certain resemblance to the doryline *Eciton*. (Freely translated from the Spanish.) Figure of mouth parts and photographs of larva in side and ventral views, p. 114.

LITERATURE CITED

- Bernard, F.** 1951. Super-famille des Formicoidea, p. 997-1104. In P. P. Grassé [ed.] *Traité de Zoologie* (Paris: Masson et Cie). Tome X, Fasc. II.
- Bruch, C.** 1925. Macho, larva y ninfa de *Acanthostichus ramosmexiae* Bruch. *Physis* 8: 110-115.
- Creighton, W. S.** 1950. The ants of North America. *Bull. Mus. Comp. Zool. Harvard College* 104: 1-585, 57 pl.
- Emery, C.** 1899. Intorno alle larve di alcune formiche. *Mem. R. Accad. Sci. Ist. Bologna* 8: 3-10.
- . 1901. Studi sul polimorfismo e la metamorfosi nel genere *Dorylus*. *Rend. Sess. R. Accad. Sc. Ist. Bologna, N. S.* 5: 415-433.
- . 1904. Le affinità del genere *Leptanilla* e i limiti delle *Dorylinae*. *Arch. Zool.* 2: 107-116.
- Kusnezov, N.** 1962. El género *Acanthostichus* Mayr. *Acta Zool. Lilloana* 18: 121-138.
- Wheeler, G. C.** 1950. Ant larvae of the subfamily *Cerapachyinae*. *Psyche* 57: 102-113.
- & **Jeanette Wheeler.** 1960. The ant larvae of the subfamily *Myrmicinae*. *Ann. Entom. Soc. Amer.* 53: 98-110.