

acterized on morphological and behavioral characters. The behavioral distinction between *Raptiformica* and such species as *F. manni*, *F. bradleyi* and their relatives have not been emphasized to the fullest extent. *Raptiformica*, as interpreted and characterized by Buren, contains only dulotic species. Those species which have been excluded from *Raptiformica* do not, so far as currently available data would indicate, ever take other species as slaves. Indeed, two of these excluded species may function as slaves to *Raptiformica*. I believe that this still further justifies the exclusion of these species from *Raptiformica*.

REFERENCES

- Buren, W. F.** 1968. Some fundamental taxonomic problems in *Formica*. Jour. Georgia Ent. Soc. 3:25-40.
- Cole, A. C., Jr.** 1946. A description of *Formica parcipappa*, a new ant from Idaho. Ann. Ent. Soc. Amer. 39:616-618.
- Creighton, W. S.** 1935. Two new species of *Formica* from western United States. Amer. Mus. Novit. 773:1-8.
- . 1950. The ants of North America. Bull. Mus. Comp. Zool. 104: 1-585.
- Gregg, R. E.** 1963. The ants of Colorado. Univ. Colo. Press, Boulder, 1-792.
- Wilson, E. O. and Brown, W. L., Jr.** 1955. Revisionary notes on the *sanguinea* and *neogagates* groups of the ant genus *Formica*. Psyche 62:108-129.