epistoma and posterior part of the abdomen somewhat hairy, the rest smooth and shining; legs long and slender; otherwise like the female.

Winged females caught about the middle of April. Dwell beneath stones and logs, having cells a few inches beneath the surface of the ground.

Hab.—Central Texas, where it is not very common, being rarely seen in the open air.

26. Formica San Sabeana, n. sp.

Female. Length 0.62 inch.—Wings extend but a little beyond the abdomen, and having one marginal and two submarginal cells, discoidal cells absent. Color: head and thorax black; abdomen yellowishbrown, the upper surface of the three last segments brownish-black; legs chestnut or yellowish-red; head subtriangular, depressed in front, vertex slightly convex; occiput emarginate, with the posterior angles rounded; eyes small, circular, and placed a little behind the middle of head on the upper margins of its sides; mandibles flattened, curved inwards, widened anteriorly, each with five teeth on their apical margins; antennæ filiform, and not enlarged towards the apical joints; prothorax large and of nearly equal width to the head, rounded above and raised higher than the mesothorax, a small depression between the meso- and metathorax; pedicle short; scale large, vertical and wedge-shaped; abdomen oblong-ovate; legs slender and short; whole ant smooth and shining, sprinkled with a very few short white hairs.

Male. Length 0.32 inch.—Abdomen very slender and oblong-ovate; color black, segments of abdomen hyaline; legs dark-brown or brownish-black; otherwise like the female. Caught in the winged state about the 1st of October.

Worker. Length 0.46 inch. Head black or brownish-black, triangular, about twice the width of the thorax; thorax yellowish-brown or piceous on its upper surface, its divisions strongly marked; otherwise like the female.

Found in an old decaying stump, in which it had many cells with intervening passages. Is very quick in its movements. Habits little known.

Hab.—Burnet and San Saba Counties, Texas. Rare.

27. Formica fœtida, n. sp.

Female. Length 0.19 inch.—Smooth throughout; honey-yellow, with brownish bands on the segments of the abdomen; head small, subtriangular, curved above and a little rounded beneath, concave be-