ed inwards, and with three small teeth near the apex, which are not seen without a good lens; epistoma depressed; antennæ inserted in front, filiform; thorax wider than the head, rounded above, its divisions obscure; pedicle short, and inserted in the base of the abdomen a little in front of the middle; scale small, inclined forwards and wedge-shaped; abdomen large, broad-ovate; head, thorax and abdomen somewhat thickly sprinkled with short grey hairs; legs slender and of medium length.

Male. Length 0.08 inch.—Head a little wider than the thorax; abdomen evate, small; otherwise like the female. Winged males and females captured in March.

Worker. Length 0.08 inch.—Color: upper surface of head, thorax and abdomen dark brown; under surface of the head, thorax and abdomen, also the pedicle, scale and legs pale yellow; head but little wider than the thorax; mesothorax depressed; abdomen ovate; otherwise like the female.

Hab.—Austin, Texas. Rare. Has cells in the ground near the surface. It is very active.

29. Formica (Tapinoma) Wichita, n. sp.

Length 0.10 inch.—Upper surface of head, thorax and abdomen jet-black and shining; legs, under surface of thorax, antennæ and mouth pale yellow; head broad-ovate, round above, at sides and behind, not emarginate; antennæ long, filiform, inserted at base of the clypeus; clypeus sub-curved, smooth; eyes rather large, prominent, sub-circular, lateral, placed about midway of the head; mandibles shorter than the labrum, very small, curved in and acute; under surface of the head not sinuate; prothorax one-third narrower than the head, round at sides, in front, above, and widest in the middle; mesothorax depressed, and a deep strangulation between it and the metathorax; metathorax widened and raised posteriorly, rounded above; pedicle inserted in the base of the abdomen a little in front of the middle; scale large, vertical, wedge-shaped; abdomen broad-ovate, obtuse; legs long; coxæ and trochanters large, a few scattering white hairs on the upper surface of the head and abdomen.

Hab.—Beneath stones near the Wichita River in Northern Texas.

30. Formica (Hypochira) subspinosa, n. sp.

Worker. Length 0.18 inch.—Yellowish-red; abdomen reddishbrown, smooth throughout; head sub-cordate, rounded above; occiput slightly emarginate, posterior angles rounded; eyes large, circular, placed a little in front of the middle on the upper surface of the round-