

Table 2. Comparison of *Leonomyrma spinosa* Arnoldi with *Chalepoxenus muellerianus* Menozzi and *C. kutteri* Cagniant.

Characters ♂	<i>L. spinosa</i>	<i>C. kutteri</i>	<i>C. muellerianus</i>
Frontal area	not clearly marked	not clearly marked	not clearly marked
Frontal carinae	parallel, slightly divergent	slightly divergent	slightly divergent
Mandibles	dentate, 4-5 teeth	dentate, 4-5 teeth	dentate, 5 teeth
Mayrian furrows*	indistinct	indistinct	indistinct
Mesonotum	anterior part narrowing	wider, nearly as wide as long	wider, nearly as wide as long
Epinotal spines	long, fine	lacking, or blunt projection	blunt projection
Wing venation	—	identical in all 3 sp.	—
Petiole	truncated but long, with strong ventral tooth	truncated, short, with short ventral tooth	truncated, short, with very small ventral projection
Postpetiole	wide, large, with ventral spine	smaller, with short, very acute ventral spine	smaller, without ventral projection
Body hairs	long, not sparse	mostly lacking, very sparse on head, thorax	shorter, not sparse
Tibial hairs	some erect	decumbent	erect
Sculpture	head wrinkled, thorax longitudinally wrinkled, epinotum wrinkled, postpetiole shining	head wrinkled, thorax, epinotum, petioles, gaster smooth, shining	entire body coarsely wrinkled, except gaster

*Notauli