

Formica fusca var. **neogagates** Em.

Two ♀ ♀.

Formica fusca var. **densiventris** n. subsp.—Length 4.5–5 mm.—Underside of head without erect hairs. Head dull, dark fuscous above, paler below, scape same color as cheeks, head and thorax above with sparse few hairs, flagellum darker, especially near apex, thorax dull, dark above like head, along the suture a lateral irregular stripe and metathorax posteriorly paler, coxæ, femora and tibiæ dark like thorax above, anterior tibiæ slightly paler, but faintly shining, apices, trochanters and tarsi paler, tarsi with sparse hairs, scale not broadened as in *subsericea*, fuscous. Abdomen dull sericeous, dorsal segments with sparse short hairs.

Black. Apical segments pale. Two ♀ ♀.

Type, Coll. Acad. Nat. Sciences, Phila. Type locality, Beulah, N. M., Aug. 17, 1901 (H. Skinner). Two ♀ ♀, head of Dailey Canon, June 26 (T. D. A. Cockerell).

The following is submitted by Prof. Cockerell :

List of ants collected at Beulah, Sapello Canon, N. M., May 3, 1902, by T. D. A. and W. P. Cockerell, and determined by Prof. W. M. Wheeler.

1. *Camponotus pennsylvanicus* DeGeer (small var.).
2. *Myrmica rubra* L., sub sp. *scabrinodis* (small dark var.).
3. *Myrmica rubra* L., sub sp. *sulcinodoides* Em. (dark var.).
4. *Formica sanguinea* L., var. *subnuda* Emery, (originally described from British America).
5. *Formica fusca* L., var. *subsericea*.
6. *Formica fusca* L., sub sp. *subpolita* Mayr, var. *neogagates* Em.
7. *Formica rufa* L., var.
8. *Lasius niger* var. *neoniger* Em.
9. *Leptothorax canadensis* Provancher.
10. *Tapinoma sessile* Say. (small var.).

PROCTOTRYPOIDEA.

SCELIONIDÆ.

Scelio ashmeadi n. sp.—Black (legs excepted), wings hyaline, faintly milky.

♂ Length 4 mm.—Head striate anterolaterally, above the middle the head is rugoso-punctate, with sparse appressed pubescence, joints of flagellum dull sculptured. Pronotum, mesonotum and dorsolum rugoso-punctate, pubescent like head, furrows on mesonotum distinct, originating near anterior margin, terminating on posterior margin. Metanotum distinctly angled laterally, impressed medially, posterior half of impression ridged longitudinally, sides of the