

Hydropsyche Taprobanes Hagen.  $4\frac{1}{2}$  l.

— mitis Hagen. 3 l.

## VI. HYMÉNOPTÈRES.

### *Formicides.*

*Formica smaragdina* F.

— mitis Smith.

— Taprobanes Smith.

— variegata Smith.

— exercita Walk. ♀  $3\frac{1}{2}$  l.

— exundans Walk. ♂ 4 l.

— meritans Walk. ♂ 5 l.

— ingruens Walk. ♀  $2\frac{1}{2}$  l.

— consultans Walk. ♂, 3 l.

— indeflexa Walk. ♀ 4 l.

*Formica pyrrhocephala* Motsch.

♂ *Elongata, sparsim pilosa, sericeo-opaca, cinereo-fusca, capite, antennis, tibiis tarsisque rufis, abdominis segmentis subalbo-marginatis; capite elongato, fronte subinfuscata, antice inter antennis late-elevata, utrinque carinata, ocellis minutissimis, vix distinctis, clypeo arcuato, medio elevato, antennis corporis fere longitudine, filiformibus; thorace capite angustiore, elongato, postice subarcuatim truncato; squama crassa, subquadrata; abdomine ovato, capitis magnitudine; tibiis tarsisque gracilibus, his valde elongatis.*

Long.  $2\frac{1}{2}$  l. — lat. cap.  $\frac{1}{2}$  l.

Des environs de Colombo.