

Female (length 3 lines) of a brown testaceous red; the legs and antennæ are pale testaceous, as are also the mandibles and apex of the metathorax; the scale is small, and not notched; the head and disk of the thorax very smooth and shining; the wings ample, and deeply smoky towards their base; the nervures and stigma are pale testaceous; the head is of the same width as the thorax, the eyes pubescent.

This species closely resembles *F. flava*, but the thorax is more elongate and narrower, the legs are stouter, and the eyes are pubescent; the female only has hitherto been captured in England.

FORMICA NIGRA.

Formica nigra, Linn. *Faun. Suec.* 1723. *Nyland. Adno. Mon. Form. Boreal.* 920, 16.

Formica fusca, Foerster, *Hym. Stud. Form.* 33, 14.

FORMICA FULIGINOSA.

Formica fuliginosa, Latr. *Fourm.* 140, t. 5, f. 27, ♂, ♀, ♂. *Nyland. Adno. Mon. Form. Boreal.* 915, 12. Foerster, *Hym. Stud. Form.* 28, 11.

FORMICA FLAVA.

Formica flava, Fab. *Ent. Syst.* ii. 357, 34. *Nyland. Adno. Mon. Form. Boreal.* 922, 17. Foerster, *Hym. Stud. Form.* 38, 17.

FORMICA FUSCA.

Formica fusca, Linn. *Faun. Suec.* 1722. *Nyland. Adno. Mon. Form. Boreal.* 919, 15.

MYRMICA RUBRA.

B. M.

Formica rubra, Linn. *Faun. Suec.* 1725.

Myrmica cœspitum, Zett. *Ins. Lapp.* 450, 1, ♂.

Myrmica ruginodis, Nyland. *Adno. Mon. Form. Boreal.* 930, 3.

Foerster, *Hym. Stud. Form.* 66, 36.