

7. DORYLUS HINDOSTANUS. Plate I. fig. 1.

Length 1 inch.—Rufo-fuscous, smooth, and covered with fine glittering downy pile: mandibles slender and acuminate at the apex; the face, base of the mandibles, the coxæ, and apex of the abdomen with long curling pale yellow hair; face and forehead slightly prominent; the ocelli in an equilateral triangle, the face deeply sulcated in front of the anterior one; the antennæ setaceous, the scape about half the length of the flagellum. Thorax gibbous in front, the scutellum transverse; the metathorax abruptly truncated, the margin of the truncation fringed with pale hairs; wings hyaline, nervures brown, the radial nervure extending a little beyond the inner angle of the marginal cell; the legs castaneous. Abdomen elongate and cylindrical, the peduncle quadrate-convex; compressed beneath into a subtriangular form; the peduncle much narrower than the following segment; the terminal segment very pilose.

Hab. India (Punjaub).

This species is most closely allied to the *D. Juvenculus*, Shuck., but may be distinguished by the different form of the under side of the peduncle; its forehead is less convex, and its peduncle narrower in proportion to the following segment.

SECT. 4.—*Peduncle quadrate; mandibles broad and nearly triangular.*

8. DORYLUS ORIENTALIS.

B.M.

Dorylus Orientalis, Westw. *Proc. Zool. Soc.* 1835, p. 72.

Shuck. Ann. & Mag. Nat. Hist. 1st ser. v. 320. 7 (1840).

Westw. Arc. Ent. i. 80. 7.

Hab. India (Bengal).

9. DORYLUS LONGICORNIS.

B.M.

Dorylus longicornis, *Shuck. Ann. & Mag. Nat. Hist.* 1st ser. v. 321. 8 (1840).

Westw. Arc. Ent. i. 80. 8.

Hab. India (Bengal).

10. DORYLUS ATTENUATUS.

Dorylus attenuatus, *Shuck. Ann. & Mag. Nat. Hist.* 1st ser. v. 322. 9 (1840).

Westw. Arc. Ent. i. 80. 9.

Hab. Africa (Gambia?). (Coll. F. Smith.)