

11. DORYLUS ATRICEPS.

Dorylus atriceps, *Shuck. Ann. Nat. Hist.* 1st ser. v. 323. 10 (1840).
Westw. Arc. Ent. i. 80. 10.

Hab. Africa (Gambia). (Coll. F. Smith.)

SECT. 5.—*Peduncle cup-shaped, as large as the following segment; the mandibles broad and nearly triangular.*

12. DORYLUS ATRATUS.

Length 9 lines.—Black, with a silky brown pubescence; elongate and cylindrical: head as wide as the thorax; the ocelli in a triangle on the vertex, which is very convex; antennæ short and setaceous, the scape more than half the length of the flagellum. Thorax: the mesothorax with an impressed longitudinal line on each side, two-thirds of its entire length, terminating at the base of the scutellum; wings blackish brown, with a slight violet iridescence, the nervures black; the recurrent nervure inserted a little beyond the middle of the first submarginal cell; femora elongate, triangular, much narrowed at the apex. Abdomen cylindrical; the peduncle cup-shaped, nearly as large as the second segment, its posterior margin rounded; beneath compressed into an angular form, and densely pubescent.

Hab. Old Calabar. (Coll. F. Smith.)

Genus 2. RHOGMUS.

Rhogmus, *Shuck. Ann. & Mag. Nat. Hist.* 1st ser. v. 323. 4 (1840).

1. RHOGMUS FIMBRIATUS. Plate I. fig. 2. B.M.

Rhogmus fimbriatus, *Shuck. Ann. & Mag. Nat. Hist.* 1st ser. v. 325. 1 (1840).

Westw. Arc. Ent. i. 80. 1.

Hab. Africa (Gold Coast; Gambia).

Genus 3. LABIDUS*.

Labidus, *Jurine, Hym.* p. 282 (1808).

SECT. 1.—*Peduncle of the abdomen subtriangular and concave above.*

1. LABIDUS FARGEAVII.

Labidus Fargeavii, *Shuck. Ann. Nat. Hist.* 1st ser. v. 198. 1 (1840).

* The females of this genus are not known. Mr. Shuckard has indicated their probable connexion with the genus *Typhlopone*.