

Labidus Fargeavii, *St. Farg. Hym.* i. 229, nec *Jurine*.
Westw. Arc. Ent. i. 74. 1.

Hab. Brazil (St. Catherine).

2. LABIDUS JURINII.

Labidus Jurinii, *Shuck. Ann. Nat. Hist.* 1st ser. v. 198. 2 (1840).
Westw. Arc. Ent. i. 75. 3.

Hab. South America (Demerara); Brazil.

3. LABIDUS BURCHELLII. Plate I. fig. 3.

Labidus Burchellii, *Westw. Arc. Ent.* i. 74. 2. pl. 20. f. 1.

Hab. Brazil (Santos).

4. LABIDUS LATREILLII.

Labidus Latreillii, *Jurine, Hym.* p. 282.
Shuck. Ann. & Mag. Nat. Hist. 1st ser. v. 199. 3 (1840).
Westw. Arc. Ent. i. 75. 4.

Latr. Gen. Crust. et Ins. iv. 123; *Nouv. Dict. d'Hist. Nat.*
2nd ed. xvii. 141.

Hab. Brazil; Demerara?; Cayenne.

5. LABIDUS SERVILLEI.

Labidus Servillei, *Westw. Arc. Ent.* i. 75. 5. pl. 20. f. 2.

Hab. Brazil (Para).

6. LABIDUS ATRICEPS.

Length 7 lines.—Body reddish, pubescent; head black; the face anteriorly, and the ridges between the antennæ at their base, the mandibles and antennæ, rufo-testaceous; the ocelli of glassy brightness, and placed in a triangle on the vertex. The anterior wings fulvo-hyaline, with the nervures brownish red; the thorax punctured above, not very closely so; on each side an impressed line runs from the scutellum nearly to the anterior margin of the mesothorax; the posterior tibiæ as wide as the femora, narrowed at the base, two-thirds of their length towards the apex of the same width, and narrowed from thence to the base; the basal joint of the tarsi broadest at the base. Abdomen curved; the peduncle of a subtriangular shape above, concave, with the margins raised, and smooth and shining. The insect covered with a fulvous pubescence.

Hab. Brazil (Ega, on the Amazons).

I have scarcely a doubt of this being the *Labidus Latreillii*, although in one particular it somewhat differs. In the original description of the 'Nouveau Dictionnaire d'Hist. Nat.,' the tibiæ