

9. LABIDUS PILOSUS.

B.M.

Length $9\frac{1}{2}$ lines.—Rufo-testaceous: the head blackish, excepting the anterior margin of the face, the mandibles, and antennæ, which are of the same colour as the body. The thorax of a darker colour above than the abdomen, punctured and shining; the sides, the metathorax, and the legs with long pale pubescence; the wings hyaline, the nervures ferruginous; the mesothorax not gibbous in front. Abdomen covered with dense shining silky pile; the peduncle transverse, subtriangular, with the sides slightly rounded.

Hab. Brazil (St. Paul's). (Coll. H. W. Bates.)

10. LABIDUS MEXICANUS.

Length 6 lines.—Pale rufo-testaceous, covered with pale shining pubescence: eyes large, black, and prominent; the ocelli glassy bright; on each side of the anterior one an elevated carina, which passes between the antennæ to the anterior margin of the face, forming a deep smooth channel; the antennæ shorter than the thorax, stout, and tapering to a point at their apex. Thorax very gibbous in front and at the scutellum; the wings wanting. Abdomen: the margins of the segments slightly constricted; the peduncle transverse, the lateral margins not elevated, slightly convex in the middle.

Hab. Mexico (Orizaba).

SECT. 2.—*Peduncle of the abdomen more or less convex, transverse-quadrate.*

11. LABIDUS ILLIGERI.

Labidus Illigeri, *Shuck. Ann. & Mag. Nat. Hist.* 1st ser. v. 397. *Append.* (1840).
Westw. Arc. Ent. i. 76. 8.

Hab. Brazil.

12. LABIDUS HARTIGII.

Labidus Hartigii, *Westw. Arc. Ent.* i. 75. 6. pl. 20. f. 3.

Hab. Brazil (Rio Janeiro and Sapé).

13. LABIDUS ESENBECKII.

Labidus Esenbeckii, *Westw. Arc. Ent.* i. 75. 7. pl. 20. f. 4.

Hab. Brazil (Rio Vendinha).