

weakly depressed; alitrunk with tegulae, mesopleuron with an oblique furrow; head, pronotum, and gaster dark reddish brown.

Holotype: Worker, Sanchahe, 960 m, Mengyang Town, Jinghong County, Yunnan Province, Feb. 27, 1997, No. A97-54 (Tiansheng Li). Paratype: 1 dealate female, with same data as holotype, No. A97-51 (Zhenghui Xu).

This new species is close to *E. buttelreepeni* Forel (Figs. 3-4), but body in colour black; in full face view occipital margin deeply concave; posteroventral angle of subpetiolar process without a denticle.

2 *Myopias* Roger

Myopias Roger 1861 *Berl. ent. Zeit.* 5:39. Type-species: *M. amblyops* Roger.

Systematic position: Ponerinae, Ponerini.

Myopias conicara sp. nov. (Figs. 5-7)

Holotype worker: TL 7.5, HL 1.50, HW 1.40, CI 93, SL 1.33, SI 96, PW 1.00, AL 2.20, ED 0.25, ML 1.35, MI 90, PNL 0.37. Head slightly longer than broad, broadest in front, narrowed posteriorly; in full face view, occipital margin nearly straight, occipital corners roundly prominent. Eyes placed on lateral sides, close to the bases of mandibles. Mandibles long and slender, slightly curved inward; inner margin with a blunt tooth near the center; masticatory margin narrow, oblique and slightly concave, with a very small denticle at basal 1/3, apical tooth acute, basal tooth triangular. Middle part of clypeus triangular, anterior margin roundly concave, anterolateral corners protruding and dentiform. Frontal lobes large, laterally expanded. Frontal furrow reached to midline of head. Antennal scapes surpass occipital corners by about 1/8 of its length; flagella incrassate towards apex, apical segment slightly longer than the preceding 2 segments combined. Ventral face of head with a conical tubercle in the middle and close to the anterior margin. In profile view, dorsum of alitrunk at the same level, promesonotal suture distinct and depressed, meta-notal groove slightly depressed. Mesonotum short, crescent in dorsal view. Dorsum of propodeum slightly convex, rounded into declivity, the latter nearly truncate, about as long as dorsum. In profile view, petiolar node quadrate, slightly higher than long; anterior face very weakly concave; posterior face truncate, each side with a vertical narrow depression close to the lateral margin; dorsal face roundly convex, anterodorsal angle and posterodorsal angle rounded, at the same height. In dorsal view, petiolar node about as long as broad, narrowing forwards, anterior margin slightly convex, anterolateral corners rounded, posterior margin straight. Subpetiolar process with a posteroventrally pointed triangular denticle at anteroventral corner, and with a