

and legs slightly lighter; gaster dark brown; pubescence yellow; pilosity brownish yellow.

Paratypes 5 workers; TL 3.1-4.0, HL 0.87-1.00, HW 0.83-1.10, CI 95-101, SL 0.86-0.75, SI 74-81, PW 0.55-0.60, AL 1.03-1.19, ED 0.17-0.22.

Holotype worker, Beiliu City, Guangxi Province of China, Aug. 26, 1995, collected by Wenming Shen. Paratypes 5 workers, data as holotype.

This new species resembles *P. cruda* (Smith), but differs from the latter in: anterior clypeal border straight; mandible with 10 teeth, followed by a row of distinct denticles on rounded basal margin distally; erect hair moderately dense, pubescence abundant.

Acknowledgements Many thanks to Professor Steven O. Shattuck (CSIRO, Division of Entomology, Australia) for his kindly sending us taxonomic papers; thanks also to Mr. Weming Shen (Guangxi Normal University) for collecting the specimens.

References

- Mann, W. M. 1921 The ants of Fiji Islands. *Bull. Mus. Comp. Zool.* **64**: 401-499.
- Shattuck, S. O. 1992a Review of the Dolichoderine ant genus *Iridomyrmex* Mayr with descriptions of three new genera (Hymenoptera; Formicidae). *J. Aust. Entomol. Soc.* **31**(1):13-18.
- Shattuck, S. O. 1992b Generic revision of the ant subfamily Dolichoderinae (Hymenoptera; Formicidae). *Sociobiology* **21**(1): 1-181.
- Smith, F. 1859 Catalogue of hymenopterous insects collected by Mr. A. R. Wallace at the Islands of Aru and Key. *J. Proc. Linn. Soc. Zool.* **3**: 132-158.
- Smith, F. 1860 Catalogue of hymenopterous insects collected by Mr. A. R. Wallace in the Islands of Bachian, Kaisaa, Amboyna, Gilolo, and at Dory in New Guinea. *J. Proc. Linn. Soc. Zool.* **5**: 93-143.

中国蚁科新纪录属及一新种记述 (膜翅目: 蚁科)

周善义 (广西师范大学生物系, 桂林 541004)

郑哲民 (陕西师范大学动物研究所, 西安 710062)

本文记述蚁科昆虫—中国新纪录属, 凹头臭蚁属 *Philidris* Shattuck 及该属一新种: 南方凹头臭蚁 *P. notiala* sp. nov.。新种与粗凹头臭蚁 *P. cruda* (Smith) 相似, 主要区别为: 唇基前缘平直; 上颚咀嚼缘具10齿, 其基缘远端具一系列清晰的细齿; 体具中等密度的直立毛和丰富的茸毛。

正模工蚁, 广西北流市, 1995年8月26日, 沈文铭采。副模5工蚁, 记载同正模。正模标本保存于陕西师范大学动物研究所, 副模标本保存于广西师范大学生物系。