



Figures 7-12. *Camponotus kaura*. 7. Major worker, frontal view of head. 8. Minor worker, same. 9. Gyne, same. 10. Major worker, lateral view of mesosoma. 11. Gyne, same. 12. Major worker, mandibles and lower clypeus, enlarged ca. 2 $\times$ , slightly oblique view. Figures 13-16. *C. taino*. 13. Major worker, frontal view of head. 14. Minor worker, same. 15. Major worker, lateral view of mesosoma. 16. Major worker, mandibles and lower clypeus, enlarged ca. 1.2 $\times$ , slightly oblique view. Figures 7-11 to same scale; Figures 13-15 to same scale.

diameter of antennal scape. Frontal lobes broad, greatest intercarinal distance about  $0.4 \times \text{HW}$ ; upper intercarinal distance about  $0.9 \times$  greatest intercarinal distance. Clypeal midline subangular for most of its length; disc slightly depressed on either side above lower margin; free margin shallowly concave and with blunt median tooth (Fig. 12); in profile, evenly sloping to ventral margin.

Antennal scape distinctly widened distad, extending to, or nearly to, level of dorsolateral lobe. Mandible with 6 teeth.

Front of head slightly to moderately shiny, surfaces finely tessellate; entire front of head sparsely and minutely punctate, clypeus least obviously punctate; mandible moderately shiny between sparse fine punctures that become finer, closer, and